Antineutrophil cytoplasmic antibody-positive conversion and microscopic polyangiitis development in patients with idiopathic pulmonary fibrosis

Naho Kagiya,¹ Noboru Takayanagi,¹ Tetsu Kanauchi,² Takashi Ishiguro,¹ Tsutomu Yanagisawa,¹ Yutaka Sugita¹

To cite: Kagiya N, Takayanagi N, Kanauchi T, *et al.* Antineutrophil cytoplasmic antibody-positive conversion and microscopic polyangiitis development in patients with idiopathic pulmonary fibrosis. *BMJ Open Res* 2015;**2**: e000058. doi:10.1136/bmjresp-2014-000058

Received 6 August 2014
Revised 22 November 2014
Accepted 1 December 2014

CrossMark

¹Department of Respiratory Medicine, Saitama Cardiovascular and Respiratory Center, Saitama, Japan

²Department of Radiology, Saitama Cardiovascular and Respiratory Center, Saitama, Japan

Correspondence to
Dr Naho Kagiya;
kagiya.naho@pref.saitama.lg.jp

ABSTRACT

Background: Increasing evidence indicates that antineutrophil cytoplasmic antibody (ANCA)-positive conversion occurs in patients initially diagnosed with idiopathic pulmonary fibrosis (IPF) and as a result, some of these patients develop microscopic polyangiitis (MPA). However, the incidence density of these patients is not well known.

Objectives: To explore the incidence of ANCA-positive conversion and development of MPA during the disease course in patients with IPF and to evaluate whether corticosteroid therapy reduces MPA development in patients with IPF with myeloperoxidase (MPO)-ANCA positivity at diagnosis or who later acquire MPO-ANCA positivity.

Methods: We retrospectively analysed the medical records of 504 Asian patients with IPF treated at our institution in Saitama, Japan.

Results: Of the 504 patients with IPF, 20 (4.0%) had MPO-ANCA and 16 (3.2%) had PR-3-ANCA when first evaluated. In 264 of 504 patients with IPF, ANCA was measured repeatedly and seroconversion to MPO-ANCA and PR3-ANCA occurred in 15 (5.7%) and 14 (5.3%) patients, respectively, and 9 of 35 patients who were either MPO-ANCA positive at IPF diagnosis or who subsequently seroconverted developed MPA. None of the nine patients who developed MPA had been previously treated with steroids. The incidence of MPA tended to be lower in patients treated than not treated with corticosteroids although this was not statistically significant.

Conclusions: Some patients with IPF with MPO-ANCA positivity at IPF diagnosis or with MPO-ANCA-positive conversion during follow-up developed MPA. Clinical trials to determine whether corticosteroid therapy can reduce MPA development and prolong survival in MPO-ANCA-positive patients with IPF should be considered.

INTRODUCTION

Vasculitides associated with serum positivity for antineutrophil cytoplasmic antibodies (ANCAs) are commonly recognised as

KEY MESSAGES

- ▶ Antineutrophil cytoplasmic antibody (ANCA)-positive conversion occurs in patients initially diagnosed with idiopathic pulmonary fibrosis (IPF) and as a result, some of these patients develop microscopic polyangiitis (MPA). However, the incidence density of these patients is not well known.
- ▶ Some patients with MPO-ANCA positivity at IPF diagnosis or with MPO-ANCA-positive conversion during follow-up developed MPA.
- ▶ The incidence of MPA tended to be lower in patients treated than not treated with corticosteroids although this was not statistically significant.

ANCA-associated vasculitis.^{1 2} ANCAs directed against proteinase 3 (PR3) are detected mainly in patients with granulomatosis with polyangiitis, whereas ANCAs directed against myeloperoxidase (MPO) are found predominantly in patients with microscopic polyangiitis (MPA) and eosinophilic granulomatosis with polyangiitis.³ Prevalences of MPO-ANCA and PR3-ANCA positivity in patients with MPA are reported to be 30–80% and 10–30%, respectively.⁴ MPA with pulmonary involvement is seen in up to 30% of patients⁵ in whom diffuse alveolar haemorrhage with pathological capillaritis is the most common manifestation.

Nada *et al*⁶ reported three patients initially diagnosed with idiopathic pulmonary fibrosis (IPF) who developed pulmonary-renal vasculitis. An association between MPA and pulmonary fibrosis has since been demonstrated.^{7–25} There sometimes appears to be an association between MPO-ANCA, MPA and pulmonary fibrosis.^{15 26} Pulmonary fibrosis associated with MPO-ANCA or MPA is either idiopathic or

associated with connective tissue disease,²⁷ and the most frequent histological or radiological pattern is that of usual interstitial pneumonia.^{15–17 22–27} MPO-ANCA-positive IPF is recognised as a distinct phenotype of IPF.²⁸

At least two possibilities have been proposed for the development of pulmonary fibrosis in patients with MPO-ANCA or MPA: repeated episodes of alveolar haemorrhage due to pulmonary capillaritis could be the pathogenesis of pulmonary fibrosis,¹⁰ and MPO-ANCAs may play a direct role in the pathogenesis of pulmonary fibrosis.^{16 29} A third hypothesis, proposed by Tzelepis *et al*,¹⁷ states that because pulmonary fibrosis is clinically manifest at MPA diagnosis, the possibility of IPF inducing MPA cannot be entirely excluded. Thereafter, Ando *et al*²³ reported that during the disease course of 61 patients with IPF, MPO-ANCA positive conversion occurred in 6 patients, of whom 2 were complicated by MPA. This implies that some ANCA-negative patients with IPF acquire ANCA positivity and then develop MPA. Taken together, these findings imply that pulmonary fibrosis is not only a consequence of MPA or of a direct role of MPO-ANCA but also that it may induce ANCA and MPA.

We thus thought that the prevalence of MPO-ANCA or PR3-ANCA positivity at IPF diagnosis and the incidence density of MPO-ANCA or PR3-ANCA-positive conversion and MPA development during follow-up should be elucidated on a large scale. Therefore, we investigated risk

factors of ANCA-positive conversion and whether ANCA positivity at IPF diagnosis was associated with mortality, and evaluated whether corticosteroid therapy reduces MPA development and prolongs survival in patients with IPF with MPO-ANCA positivity at diagnosis or who later acquire MPO-ANCA positivity.

METHODS

Participants

From 1998 through 2012, 966 patients with IPF were treated at our institution (figure 1). Of these patients, 462 were excluded: 10 patients with MPA at IPF diagnosis, 21 with exacerbation of IPF at diagnosis, 248 with simultaneous lung cancer, 39 with simultaneous chronic pulmonary infections and 144 patients in whom ANCA measurement was unavailable. Patients with IPF and lung cancer or chronic pulmonary infections were not included as these conditions can be associated with ANCA positivity.³⁰ Thus, 504 patients comprised the cohort of this study, and they were further divided into three cohorts according to ANCA positivity at IPF diagnosis whether ANCA had or had not been measured repeatedly (figure 1). Patients were followed up through August 2013 or until death. All patients fulfilled the criteria for IPF of the American Thoracic Society and European Respiratory Society³¹ or the official ATS/ERS/

Figure 1 Flow diagram of enrolment and median follow-up periods in patients with idiopathic pulmonary fibrosis (IPF), ANCA, antineutrophil cytoplasmic antibody; MPO, myeloperoxidase; PR3, proteinase 3.

Table 1 Baseline characteristics of the study patients with IPF according to ANCA positivity at diagnosis

Characteristic	Total	ANCA		p Value*
		Positive	Negative	
Number of patients	504	20	468	
Male	376 (74.6%)	11 (55.0%)	354 (75.6%)	0.072
Age, years	69.5±8.1	71.4±7.6	69.3±8.2	0.041
Smoker	383 (76.0%)	11 (55%)	359 (76.7%)	0.414
Emphysema	131 (26.0%)	5 (25.0%)	122 (26.1%)	1
%FVC predicted	74.7±19.7	68.9±23.9	75.3±19.3	0.011
FEV1/FVC, %	81.1±10.7	81.1±14.8	80.9±10.6	0.716
DLCO, % predicted	77.0±23.4	74.0±11.5	77.5±23.6	0.124
WCC, /μL	7292±2213	8440±3614	7227±2139	0.078
ESR, mm/h	37.8±27.9	74.1±41.5	35.8±26.1	<0.001
Creatinine, mg/dL	0.80±0.48	0.73±0.15	0.81±0.50	0.078
CRP, mg/d†	0.20 (0.10–0.63)	1.17 (0.50–5.30)	0.20 (0.10–0.60)	<0.001
KL-6, IU/L†	778 (512–1263)	646 (389–878)	784 (519–1247)	0.589
RF positive	84 (16.7%)	14 (70.0%)	68 (14.5%)	<0.001
ANA positive	284 (56.3%)	17 (85.0%)	255 (54.5%)	0.016
Urinary blood positive	35 (6.9%)	6 (30.0%)	28 (6.0%)	0.143
Urinary protein positive	30 (6.0%)	3 (15.0%)	27 (5.8%)	1

*p Values were calculated in relation to ANCA positivity.

†Median value with IQR in parentheses.

ANA, antinuclear antibody; ANCA, antineutrophil cytoplasmic antibody; CRP, C reactive protein; DLCO, diffusing capacity for carbon monoxide; ESR, erythrocyte sedimentation rate; FEV1/FVC, forced expiratory volume in 1 s/FVC ratio; FVC, forced vital capacity; IPF, idiopathic pulmonary fibrosis; KL-6, Krebs von den Lungen-6; MPO, myeloperoxidase; PR3, proteinase 3; RF, rheumatoid factor; WCC, white cell count.

JRS/ALAT statement on IPF.³² MPA was diagnosed using the Chapel Hill consensus criteria.¹ This study was approved by the institutional review board of Saitama Cardiovascular and Respiratory Center.

Study design

This was a retrospective cohort study. Clinical, radiographic, laboratory data and outcome were collected from medical records. Baseline clinical parameters were obtained within 1 month of the initial diagnosis. If these data were not obtained within this period, we considered them to be unknown. Survival status was obtained from medical records and/or telephone interviews.

Measurement of ANCA

MPO-ANCAs and PR3-ANCAs were tested using an enzyme immunoassay: January 1998–August 1999 (NIPRO Japan), September 1999–March 2002 (BIO-RAD, USA), April 2002–March 2012 (Medical and Biological Laboratories, Japan) and April 2012 onward (Phadia Laboratory Systems, Japan).

Statistical analysis

Categorical baseline characteristics are summarised by frequency and per cent, and continuous characteristics are reported as the mean±SD or median and IQR as appropriate. Group comparisons were made using a

Figure 2 Kaplan–Meier curves for the time until myeloperoxidase-antineutrophil cytoplasmic antibody (MPO-ANCA)-positive conversion (A: MPO-ANCA, B: proteinase 3 [PR3]-ANCA) in patients with idiopathic pulmonary fibrosis.

student *t* test, Wilcoxon rank-sum tests, or Fisher's exact test, as appropriate. ANCA-positive conversion and MPA diagnosis were estimated by Kaplan–Meier analysis. Survival was evaluated using a Kaplan–Meier curve and compared between groups using log-lank tests. Cox regression analysis was used to determine whether the following factors increased the risk of ANCA-positive conversion: sex, age, smoking history, emphysema, forced vital capacity (FVC), forced expiratory volume in 1 s (FEV1)/FVC ratio, lung diffusion capacity for carbon monoxide (DLCO), white cell count, erythrocyte sedimentation rate (ESR), C reactive protein, serum creatinine, urine protein, urine blood, rheumatoid factor, antinuclear antibody and Krebs von den Lungen-6 (KL-6). Rheumatoid factor ≥ 20 IU/mL and antinuclear antibody $\geq 1/80$ were considered to indicate positivity. Emphysema was considered present if areas of low attenuation were present on high-resolution CT images. Cox regression analysis was used to determine whether the following factors increased patient mortality: sex, age, smoking history, emphysema, FVC, FEV1/FVC, DLCO, MPO- or PR3-ANCA positivity, white cell count, ESR, C reactive protein, serum creatinine, urine protein, urine blood, rheumatoid factor, antinuclear antibody and KL-6. In all analyses, a *p* value of <0.05 was considered to be statistically significant. We conducted all statistical analyses with SAS V.9.2 (SAS Institute, Cary, North Carolina, USA).

RESULTS

Patient characteristics and ANCA positivity at IPF diagnosis

Of the 504 patients, 20 (4%) were MPO-ANCA positive and 16 (3.2%) were PR3-ANCA positive (table 1) at diagnoses. All of these patients were Asian. The ANCA-positive patients were older than the ANCA-negative patients. ESR and C reactive protein values were higher and FVC was lower in the ANCA-positive versus ANCA-negative patients. Rheumatoid factor positivity and antinuclear antibody positivity were more likely to be seen in the ANCA-positive patients than in the ANCA-negative patients.

Incidence density of ANCA-positive conversion

During the disease course, MPO-ANCA or PR3-ANCA-positive conversion occurred in 15 (5.7%) and 14 (5.3%), respectively, of the 264 patients in whom ANCA was repeatedly measured over a median follow-up period of 5.03 years (IQR, 3.11–8.07 years). Thus, the incidence density of MPO-ANCA and PR3-ANCA-positive conversion was 13.10 and 12.23 cases per 1000 person-years, respectively (figure 2).

Risk factors for MPO- or PR3-ANCA-positive conversion

In a multivariate Cox regression hazard model, rheumatoid factor positivity was associated with MPO-ANCA-positive conversion (adjusted HR 3.435, 95% CI 1.032 to 11.440,

Table 2 Clinical features of the 19 patients with IPF and MPA

Case	Sex	Age at the time of MPA development (years)	Time from IPF diagnosis to MPA development (years)	MPO-ANCA at the time of IPF diagnosis	MPO-ANCA at the time of MPA development	Diffuse alveolar haemorrhage	Acute respiratory failure	Rapidly progressive glomerulonephritis	Mononeuritis multiplex	Gastrointestinal bleeding	Purpuric rash	Fever	Duration after MPA development (years)	Outcome
1	M	67	0	Positive	Positive	+	+	+	+	+	+	+	9.04	Dead
2	M	67	0	Positive	Positive	+	+	+	+	+	+	+	2.41	Dead
3	F	72	0	Positive	Positive	+	+	+	+	+	+	+	0.27	Alive
4	F	66	0	Positive	Positive	+	+	+	+	+	+	+	0.6	Dead
5	F	73	0	Positive	Positive	+	+	+	+	+	+	+	0.16	Alive
6	M	62	0	Positive	Positive	+	+	+	+	+	+	+	9.99	Alive
7	M	69	0	Positive	Positive	+	+	+	+	+	+	+	3.69	Alive
8	F	79	0	Positive	Positive	+	+	+	+	+	+	+	1.53	Alive
9	M	69	0	Positive	Positive	+	+	+	+	+	+	+	2.47	Alive
10	F	79	0	Positive	Positive	+	+	+	+	+	+	+	0.08	Dead
11	F	74	0.28	Positive	Positive	+	+	+	+	+	+	+	1.32	Dead
12	M	75	6.06	Negative	Positive	+	+	+	+	+	+	+	0.06	Dead
13	M	76	5.5	Negative	Positive	+	+	+	+	+	+	+	0.08	Dead
14	M	76	5.12	Negative	Positive	+	+	+	+	+	+	+	1.04	Dead
15	F	73	8.25	Negative	Positive	+	+	+	+	+	+	+	1.87	Dead
16	M	69	0.5	Positive	Positive	+	+	+	+	+	+	+	9.76	Alive
17	F	69	6.93	Positive	Positive	+	+	+	+	+	+	+	0.08	Alive
18	F	60	0.42	Negative	Positive	+	+	+	+	+	+	+	6.31	Alive
19	M	62	5.33	Negative	Positive	+	+	+	+	+	+	+	0.08	Alive

ANCA, antineutrophil cytoplasmic antibody; IPF, idiopathic pulmonary fibrosis; MPA, microscopic polyangiitis.

Figure 3 Kaplan–Meier curves for the time until development of microscopic polyangiitis (MPA) in patients with idiopathic pulmonary fibrosis according to antineutrophil cytoplasmic antibody (ANCA) positivity at diagnosis. The log-rank test showed the difference between ANCA-negative patients and myeloperoxidase (MPO)-ANCA-positive patients to be significant ($p<0.001$). PR3, proteinase 3.

$p=0.044$), as was an ESR of ≥ 40 mm/h (adjusted HR 3.361, 95% CI 1.100 to 10.271, $p=0.033$).

Incidence density of MPA development according to ANCA positivity

Ten patients with MPA at IPF diagnosis were excluded from this analysis. Among these 10 patients (5 men, 5 women) with MPA at IPF diagnosis, the median age was 69 (range, 62–79) years, 7 had rapidly progressive glomerulonephritis, 10 had diffuse alveolar haemorrhage, 3 had acute respiratory failure, 4 had fever, 5 had mononeuritis multiplex and 2 had purpuric rash. In total, 9 patients in the cohort of 504 developed MPA. Among these 9 patients

Figure 4 Kaplan–Meier survival curves of all-cause mortality according to antineutrophil cytoplasmic antibody (ANCA) positivity. The log-rank test showed the difference between ANCA-positive and ANCA-negative survival curves to be significant ($p<0.001$).

(5 men, 4 women), the median age was 73 (range, 60–76) years, 7 had rapidly progressive glomerulonephritis, 3 had diffuse alveolar haemorrhage, 3 had acute respiratory failure, 8 had fever, and 1 patient each had mononeuritis multiplex, gastrointestinal bleeding and purpuric rash (table 2). Three of the 20 patients with MPO positivity and none of the 16 patients with PR3-ANCA positivity at IPF diagnosis developed MPA over a median follow-up period of 2.42 years (IQR 1.38–4.92 years) (incidence density of 39.4 cases per 1000 MPO-ANCA-positive IPF person-years and 0.00 cases per 1000 PR3-ANCA-positive IPF person-years, respectively; figure 3). Of the 468 patients with ANCA negativity at IPF diagnosis, 6 developed MPA over a median follow-up period of 4.02 years (IQR 1.92–6.73 years) (incidence density of 2.75 cases per 1000 ANCA-negative IPF person-years), MPO-ANCA converted to positive in 5 patients at MPA diagnosis and 1 patient developed MPA 19 months after ANCA converted to positive. MPO-ANCA-positive patients developed MPA more frequently than did ANCA-negative patients ($p<0.001$).

Mortality according to ANCA positivity at IPF diagnosis

Of the 504 patients with IPF, death from any cause occurred in 245 (48.6%) patients over a median follow-up period of 3.95 years (IQR, 1.84–6.64 years). Patients died from progression of the IPF (35.9%), acute exacerbation of IPF (21.0%), pneumonia (10.1%), lung cancer (9.3%), other pulmonary diseases (2.4%), MPA (1.2%), non-pulmonary diseases (10.9%) and unknown causes (9.2%). Five-year and 10-year mortality rates were, respectively, 61.3% and 85.7% for ANCA-positive patients, and 37.6% and 70.5% for ANCA-negative patients at IPF diagnosis. The log-rank test showed the difference between survival curves of ANCA-positive and ANCA-negative patients to be significant ($p=0.001$) (figure 4). Five-year mortality rates of MPO-ANCA and PR3-ANCA positive patients were 51.3% and 81.7%, respectively. In a multivariate Cox proportional hazard model, older age, PR3-ANCA positivity, %FVC predicted $<70\%$, FEV1/FVC $\geq 70\%$ and DLCO $<70\%$ were found to be negative prognostic factors (table 3). Since FEV1/FVC $\geq 70\%$ was an independent risk factor of IPF mortality, we compared baseline parameters of patients with IPF whose FEV1/FVC ratio was $\geq 70\%$ and $<70\%$. In comparison with patients with an FEV1/FVC $\geq 70\%$, patients with an FEV1/FVC $<70\%$ were significantly more frequently male (90.2% vs 72.5%, $p=0.006$), smokers (94.1% vs 75.3%, $p<0.001$), had emphysema (70.6% vs 21.7%, $p<0.001$) and had higher %FVC predicted (89.00 ± 18.43 vs 72.74 ± 18.97 mL, $p<0.001$).

MPA development and mortality in patients with positive and positively-converted MPO-ANCA according to corticosteroid therapy before MPA onset

In this cohort of patients with IPF, a total of 35 patients were positive for MPO-ANCA (20 at IPF diagnosis and 15 subsequent seroconversion). Among this retrospective cohort of patients with IPF from 1998 to 2013, some

Table 3 Univariate and multivariate analysis of the risk of all-cause mortality

Variable	Univariate cox regression			Multivariate cox regression final model		
	Crude HR	95% CI	p Value	Adjusted HR	95% CI	p Value
ANCA						
Negative	Reference	–	–	Reference	–	–
MPO-ANCA positive	1.647	0.959 to 2.829	0.071	1.480	0.836 to 2.619	0.178
PR3-ANCA positive	2.987	1.523 to 5.859	0.001	2.415	1.225 to 4.761	0.011
Sex						
Female	Reference	–	–			
Male	0.836	0.625 to 1.118	0.226			
Age						
<65 years	Reference	–	–	Reference	–	–
≥65 years	1.757	1.307 to 2.363	<0.001	1.694	1.253 to 2.292	<0.001
Smoking status						
Never smoker	Reference	–	–			
Ex/current smoker	0.660	0.492 to 0.887	0.006			
Emphysema						
None	Reference	–	–			
Some	0.855	0.647 to 1.130	0.272			
%FVC predicted						
≥70%	Reference	–	–	Reference	–	–
<70%	2.961	2.214 to 3.958	<0.001	2.457	1.810 to 3.334	<0.001
Unknown	1.830	1.312 to 2.552	<0.001	0.060	0.007 to 0.532	0.012
FEV1/FVC, %						
≥70%	Reference	–	–	Reference	–	–
<70%	0.526	0.331 to 0.838	0.007	0.560	0.347 to 0.920	0.017
Unknown	1.173	0.865 to 1.592	0.305	18.379	2.137 to 158.081	0.008
DLCO, % predicted						
≥70%	Reference	–	–	Reference	–	–
<70%	1.920	1.344 to 2.743	<0.001	1.586	1.099 to 2.291	0.014
Unknown	2.125	1.570 to 2.876	<0.001	1.954	1.377 to 2.773	<0.001
WCC						
<10 000/μL	Reference	–	–			
≥10 000/μL	1.657	1.157 to 2.373	0.006			
Unknown	0.352	0.111 to 1.121	0.077			
ESR						
<40 mm/h	Reference	–	–			
≥40 mm/h	1.627	1.202 to 2.203	0.002			
Unknown	1.289	0.949 to 1.751	0.104			
Creatinine						
<1.0 mg/dL	Reference	–	–			
≥1.0 mg/dL	1.178	0.838 to 1.656	0.345			
Unknown	0.701	0.326 to 1.511	0.365			
CRP						
<1.0 mg/dL	Reference	–	–			
≥1.0 mg/dL	1.243	0.916 to 1.688	0.163			
Unknown	0.542	0.276 to 1.065	0.076			
KL-6						
<1000 IU/L	Reference	–	–			
≥1000 IU/L	1.929	1.426 to 2.610	<0.001			
Unknown	0.836	0.604 to 1.158	0.282			
ANA						
Negative	Reference	–	–			
Positive	1.313	0.979 to 1.761	0.069			
Unknown	0.705	0.447 to 1.112	0.132			
Rheumatoid factor						
Negative	Reference	–	–			
Positive	1.156	0.831 to 1.610	0.390			
Unknown	0.649	0.454 to 0.928	0.018			

Continued

Table 3 Continued

Variable	Univariate cox regression			Multivariate cox regression final model		
	Crude HR	95% CI	p Value	Adjusted HR	95% CI	p Value
Urinary blood						
Negative	Reference	—	—			
Positive	1.067	0.674 to 1.687	0.782			
Unknown	0.737	0.565 to 0.962	0.025			
Urinary protein						
Negative	Reference	—	—			
Positive	1.066	0.610 to 1.865	0.822			
Unknown	0.734	0.566 to 0.951	0.020			

ANA, antinuclear antibody; ANCA, antineutrophil cytoplasmic antibody; CRP, C-reactive protein; DLCO, diffusing capacity for carbon monoxide; ESR, erythrocyte sedimentation rate; FEV1/FVC, forced expiratory volume in 1 s/FVC ratio; FVC, forced vital capacity; KL-6, Krebs von den Lungen-6; MPO, myeloperoxidase; PR3, proteinase 3; WCC, white cell count.

patients received corticosteroid treatment, although this is no longer recommended in the international guidance published in 2012.³² Among the 35 MPO-ANCA-positive patients, 8 had received systemic corticosteroids and none developed MPA. Among the remaining 27 MPO-ANCA-positive patients who never received steroids, 9 developed MPA. Accordingly, the incidence of developing MPA tended to be lower in patients treated with corticosteroids than in patients not treated with corticosteroids ($p=0.063$). There was no significant difference in survival between patients treated or not treated with corticosteroids ($p=0.323$) (figure 5).

DISCUSSION

This long-term longitudinal study of a large cohort of patients with IPF resulted in five important findings. First, the incidence density of ANCA-positive conversion in patients with IPF was 13.10 cases per 1000 person-years. Second, some patients with IPF with MPO-ANCA positivity at IPF diagnosis or with MPO-ANCA positive conversion during the disease course developed MPA.

Third, rheumatoid factor positivity was a risk factor for MPO-ANCA-positive conversion, and an ESR ≥ 40 mm/h was a risk factor for PR3-ANCA-positive conversion. Fourth, PR3-ANCA positivity at IPF diagnosis was an independent risk factor for mortality in patients with IPF. Fifth, corticosteroid therapy might reduce MPA development in patients with IPF with MPO-ANCA positivity or positive conversion.

The prevalence of MPO- and PR3-ANCA positivity in patients with IPF at diagnosis has been reported to vary between 4.9–32.1% and 2.2–3.7%, respectively,^{23 24 33} and these prevalences were comparable to ours. In patients with IPF with MPO-ANCA positivity at diagnosis, Nozu *et al*²⁴ reported that 4 of 17 patients developed MPA, and Kang *et al*³³ reported that 2 of 27 patients developed MPA. Ando *et al*²³ reported that two of six patients with MPO-ANCA-positive conversion developed MPA. These studies and ours suggest that not only patients with IPF who are MPO-ANCA positive at diagnosis but also those who convert to positive develop MPA.

Gaudin *et al*¹⁹ reported that MPO-ANCA-associated and PR3-ANCA-associated vasculitis were both associated

Figure 5 Kaplan–Meier curves for the time until development of microscopic polyangiitis (MPA) (A) and for survival (B) in myeloperoxidase-antineutrophil cytoplasmic antibody-positive or antibody-positively converted patients with idiopathic pulmonary fibrosis according to corticosteroid treatment. The incidence of the development of MPA tended to be lower in patients treated than not treated with corticosteroids (log-rank test: $p=0.063$). The difference in survival curves between patients treated and not treated with corticosteroids was not significant (log-rank test: $p=0.323$).

with pulmonary fibrosis. However, subsequent reports of patients with ANCA-associated vasculitis and IPF were restricted to MPO-ANCA-positive patients.^{15–17 22 24} In the present study, some patients with IPF had MPO-ANCA or PR3-ANCA positivity or positive conversion. However, all of the patients who developed MPA were MPO-ANCA-positive patients. Previous studies and ours indicate that MPO-ANCA should be measured regularly in patients with IPF. Although the clinical benefit of testing for PR3-ANCA might be limited, it may be helpful because our study indicated that PR3-ANCA positivity was an independent risk factor for mortality.

Arulkumaran *et al*¹⁵ reported that of 14 patients with MPA and interstitial lung disease, 2 were diagnosed as having interstitial lung disease 6 months and 3 years, respectively, before MPA. Three of their patients presented with MPA before interstitial lung disease with a mean interval of 9.7 years, and 9 patients presented with features of interstitial lung disease and MPA concurrently. If pulmonary fibrosis develops after the onset of MPA, repeated episodes of alveolar haemorrhage due to pulmonary capillaritis could be the pathogenesis of interstitial fibrosis. Tanaka *et al*²⁵ reported that in nine patients with interstitial pneumonia associated with MPO-ANCA but without MPA, surgical lung biopsy showed neither capillaritis nor vasculitis. Therefore, if patients with IPF are proved to have MPO-ANCA at IPF diagnosis, two possible pathogeneses of pulmonary fibrosis may be present: MPO-ANCA is playing a direct role in the pathogenesis of pulmonary fibrosis, and pulmonary fibrosis is inducing MPO-ANCA. During the disease course, if MPO-ANCA-positive conversion occurs, IPF might be inducing MPO-ANCA and MPA.

Therapy with corticosteroids in combination with other immunosuppressants is the current mainstay of MPA treatment.^{4 34} However, the 2012 official ATS/ERS/JRS/ALAT statement recommends that patients with IPF should not be treated with corticosteroid monotherapy or combined corticosteroid and immunomodulator therapy.³² Moreover, increased risks of death and hospitalisation were observed in patients with IPF who were treated with a combination of prednisone, azathioprine and N-acetylcysteine, as compared with placebo.³⁵ However, if patients with IPF develop MPA, patients should be treated with corticosteroids because untreated MPA is normally rapidly progressive and fatal.⁴ Thus, should patients with IPF with MPO-ANCA positivity but without MPA be treated with corticosteroids? Ando *et al* reported that 6 of 9 MPO-ANCA-positive patients with IPF were treated with corticosteroid therapy and none developed MPA, but 2 of 3 patients not treated with corticosteroids developed MPA.²³ We hypothesise that corticosteroid therapy in patients with IPF with MPO-ANCA positivity might have some benefit in reducing the development of MPA.

One limitation of this study is that it was retrospective, so some clinical and laboratory findings were not available. Second, because the decision to measure ANCA was made by referring doctors and the interval of

measurement of ANCA was not unified, we could not conclude how often ANCA should be measured in patients with IPF. Finally, because the ELISA kit had been changed four times during the study period, we could not accurately assess the association of ANCA titres with MPA development.

In conclusion, the present study showed that the incidence density of ANCA-positive conversion in patients with IPF was 13.10 cases per 1000 person-years. Some patients with IPF with MPO-ANCA positivity at IPF diagnosis or with MPO-ANCA-positive conversion during the disease course developed MPA. PR3-ANCA positivity at IPF diagnosis was found to be an independent risk factor for mortality. Clinical trials to determine whether corticosteroid therapy can reduce MPA development and prolong survival in MPO-ANCA-positive patients with IPF should be considered

Acknowledgements The authors offer their sincerest thanks to Drs Youtaro Takaku and Kazuyoshi Kurashima of the Department of Respiratory Medicine, Saitama Cardiovascular and Respiratory Center, for their handling of the diagnosis and treatment of the patients with IPF.

Contributors NK performed the primary data analysis. NT had the idea for the study. NT and TK reviewed the chest radiographs and high-resolution CT scans. NK and NT wrote the manuscript. TI, TY and YS contributed to study design, interpreted the data and reviewed the manuscript. All authors approved the final version of the manuscript.

Competing interests None.

Ethics approval This study was approved by the institutional review board of Saitama Cardiovascular and Respiratory Center.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Jennette JC, Falk RJ, Andrassy K, *et al*. Nomenclature of systemic vasculitides. Proposal of an international consensus conference. *Arthritis Rheum* 1994;37:187–92.
- Watts R, Lane S, Hanslik T, *et al*. Development and validation of a consensus methodology for the classification of the ANCA-associated vasculitides and polyarteritis nodosa for epidemiological studies. *Ann Rheum Dis* 2007;66:222–7.
- Jennette JC, Falk RJ, Bacon PA, *et al*. 2012 revised international chapel hill consensus conference nomenclature of Vasculitides. *Arthritis Rheum* 2013;65:1–11.
- Gómez-Puerta JA, Hernández-Rodríguez J, López-Soto A, *et al*. Antineutrophil cytoplasmic antibody-associated vasculitides and respiratory disease. *Chest* 2009;136:1101–11.
- Brown KK. Pulmonary vasculitis. *Proc Am Thorac Soc* 2006;3:48–57.
- Nada AK, Torres VE, Ryu JH, *et al*. Pulmonary fibrosis as an unusual clinical manifestation of a pulmonary-renal vasculitis in elderly patients. *Mayo Clin Proc* 1990;65:847–56.
- Becker-Merok A, Nossent JC, Rittland N. Fibrosing alveolitis predating microscopic polyangiitis. *Scand J Rheumatol* 1999;28:254–6.
- Hiromura K, Nojima Y, Kitahara T, *et al*. Four cases of anti-myeloperoxidase antibody-related rapidly progressive glomerulonephritis during the course of idiopathic pulmonary fibrosis. *Clin Nephrol* 2000;53:384–9.

9. Mansi IA, Opran A, Sondhi D, *et al*. Microscopic polyangiitis presenting as idiopathic pulmonary fibrosis: is anti-neutrophilic cytoplasmic antibody testing indicated? *Am J Med Sci* 2001;321:201–2.
10. Eschun GM, Mink SN, Sharma S. Pulmonary interstitial fibrosis as a presenting manifestation in perinuclear antineutrophilic cytoplasmic antibody microscopic polyangiitis. *Chest* 2003;123:297–301.
11. Birnbaum J, Danoff S, Askin FB, *et al*. Microscopic polyangiitis presenting as a “pulmonary-muscle” syndrome: is subclinical alveolar hemorrhage the mechanism of pulmonary fibrosis? *Arthritis Rheum* 2007;56:2065–71.
12. Bhanji A, Karim M. Pulmonary fibrosis-an uncommon manifestation of anti-myeloperoxidase-positive systemic vasculitis? *NDT Plus* 2010;3:351–3.
13. Shields O, Shah A, Mann B. Pyrexia of unknown origin and pulmonary fibrosis as a presentation of MPO-ANCA associated vasculitis. *BMJ Case Rep* 2011;2011. doi:10.1136/bcr.01.2011.3692
14. Eleftheriou D, Katsenos S, Zorbas S, *et al*. Pulmonary fibrosis presenting as an early manifestation of microscopic polyangiitis. *Monaldi Arch Chest Dis* 2012;77:141–4.
15. Arulkumaran N, Periselneris N, Gaskin G, *et al*. Interstitial lung disease and ANCA-associated vasculitis: a retrospective observational cohort study. *Rheumatology (Oxford)* 2011;50:2035–43.
16. Hervier B, Pagnoux C, Agard C, *et al*. Pulmonary fibrosis associated with ANCA-positive vasculitides. Retrospective study of 12 cases and review of the literature. *Ann Rheum Dis* 2009;68:404–7.
17. Tzelepis GE, Kokosi M, Tzioufas A, *et al*. Prevalence and outcome of pulmonary fibrosis in microscopic polyangiitis. *Eur Resp J* 2010;36:116–21.
18. Gal AA, Salinas FF, Staton GW Jr. The clinical and pathological spectrum of antineutrophil cytoplasmic autoantibody-related pulmonary disease. A comparison between perinuclear and cytoplasmic antineutrophil cytoplasmic autoantibodies. *Arch Pathol Lab Med* 1994;118:1209–14.
19. Gaudin PB, Askin FB, Falk RJ, *et al*. The pathologic spectrum of pulmonary lesions in patients with anti-neutrophil cytoplasmic autoantibodies specific for anti-proteinase 3 and anti-myeloperoxidase. *Am J Clin Pathol* 1995;104:7–16.
20. Ando Y, Okada F, Matsumoto S, *et al*. Thoracic manifestation of myeloperoxidase-antineutrophil cytoplasmic antibody (MPO-ANCA)-related disease. CT findings in 51 patients. *J Comput Assist Tomogr* 2004;28:710–16.
21. Chung MP, Yi CA, Lee HY, *et al*. Imaging of pulmonary vasculitis. *Radiology* 2010;255:322–41.
22. Foulon G, Delaval P, Valeyre D, *et al*. ANCA-associated lung fibrosis: analysis of 17 patients. *Respir Med* 2008;102:1392–8.
23. Ando M, Miyazaki E, Ishii T, *et al*. Incidence of myeloperoxidase anti-neutrophil cytoplasmic antibody positivity and microscopic polyangiitis in the course of idiopathic pulmonary fibrosis. *Respir Med* 2013;107:608–15.
24. Nozu T, Kondo M, Suzuki K, *et al*. A comparison of the clinical features of ANCA-positive and ANCA-negative idiopathic pulmonary fibrosis patients. *Respiration* 2009;77:407–15.
25. Tanaka T, Otani K, Egashira R, *et al*. Interstitial pneumonia associated with MPO-ANCA: clinicopathological features of nine patients. *Respir Med* 2012;106:1765–70.
26. Yamada H. ANCA: associated lung fibrosis. *Semin Respir Crit Care Med* 2011;32:322–7.
27. Homma S, Matsushita H, Nakata K. Pulmonary fibrosis in myeloperoxidase antineutrophil cytoplasmic antibody-associated vasculitides. *Respirology* 2004;9:190–6.
28. Poletti V, Ravaglia C, Buccioli M, *et al*. Idiopathic pulmonary fibrosis: diagnosis and prognostic evaluation. *Respiration* 2013;86:5–12.
29. Guilpain P, Chéreau C, Goulvestre C, *et al*. The oxidation induced by antimyeloperoxidase antibodies triggers fibrosis in microscopic polyangiitis. *Eur Respir J* 2011;37:1503–13.
30. Tsiveriotis K, Tsirogianni A, Pipi E, *et al*. Antineutrophil cytoplasmic antibodies testing in a large cohort of unselected greek patients. *Autoimmune Dis* 2011;2011:626495.
31. American Thoracic Society. Idiopathic pulmonary fibrosis: diagnosis and treatment. International consensus statement. American Thoracic Society (ATS), and the European Respiratory Society (ERS). *Am J Respir Crit Care Med* 2000;161:646–64.
32. Raghu G, Collard HR, Egan JJ, *et al*. An official ATS/ERS/JRS/ALAT statement: idiopathic pulmonary fibrosis: evidence-based guidelines for diagnosis and management. *Am J Respir Crit Care Med* 2011;183:788–824.
33. Kang BH, Park JK, Roh JH, *et al*. Clinical significance of serum autoantibodies in idiopathic interstitial pneumonia. *J Korean Med Sci* 2013;28:731–7.
34. Guillevin L, Durand-Gasselin B, Cevallos R, *et al*. Microscopic polyangiitis: clinical and laboratory findings in eighty-five patients. *Arthritis Rheum* 1999;42:421–30.
35. Raghu G, Anstrom KJ, King TE Jr, *et al*. Idiopathic Pulmonary Fibrosis Clinical Research Network, Prednisone, azathioprine, and N-acetylcysteine for pulmonary fibrosis. *N Engl J Med* 2012;366:1968–77.