

# Using domiciliary non-invasive ventilator data downloads to inform clinical decision-making to optimise ventilation delivery and patient compliance

Stephanie K Mansell,<sup>1,2</sup> Steven Cutts,<sup>1</sup> Isobel Hackney,<sup>1</sup> Martin J Wood,<sup>3</sup> Kevin Hawksworth,<sup>3</sup> Dean D Creer,<sup>2</sup> Cherry Kilbride,<sup>1,4</sup> Swapna Mandal<sup>2</sup>

**To cite:** Mansell SK, Cutts S, Hackney I, *et al.* Using domiciliary non-invasive ventilator data downloads to inform clinical decision-making to optimise ventilation delivery and patient compliance. *BMJ Open Resp Res* 2018;**5**:e000238. doi:10.1136/bmjresp-2017-000238

Received 9 August 2017  
Revised 14 December 2017  
Accepted 7 January 2018

## ABSTRACT

**Introduction** Ventilation parameter data from patients receiving home mechanical ventilation can be collected via secure data cards and modem technology. This can then be reviewed by clinicians and ventilator prescriptions adjusted. Typically available measures include tidal volume ( $V_T$ ), leak, respiratory rate, minute ventilation, patient triggered breaths, achieved pressures and patient compliance. This study aimed to assess the potential impact of ventilator data downloads on management of patients requiring home non-invasive ventilation (NIV).

**Methods** A longitudinal within-group design with repeated measurements was used. Baseline ventilator data were downloaded, reviewed and adjustments made to optimise ventilation. Leak,  $V_T$  and compliance data were collected for comparison at the first review and 3–7 weeks later. Ventilator data were monitored and amended remotely via a modem by a consultant physiotherapist between the first review and second appointment.

**Results** Analysis of data from 52 patients showed increased patient compliance (% days used >4 hours) from 90% to 96% ( $p=0.007$ ), increased usage from 6.53 to 6.94 hours ( $p=0.211$ ) and a change in  $V_T$  (9.4 vs 8.7 mL/kg/ideal body weight,  $p=0.022$ ). There was no change in leak following review of NIV prescriptions (mean (SD): 43 (23.4) L/min vs 45 (19.9) L/min,  $p=0.272$ ).

**Conclusion** Ventilator data downloads, via early remote assessment, can help optimise patient ventilation through identification of modifiable factors, in particular interface leak and ventilator prescriptions. However, a prospective study is required to assess whether using ventilator data downloads provides value in terms of patient outcomes and cost-effectiveness. The presented data will help to inform the design of such a study.

## INTRODUCTION

Domiciliary non-invasive ventilation (NIV) is accepted as standard care for patients with chronic hypercapnic ventilatory failure.<sup>1,2</sup> The efficacy of domiciliary NIV is dependent on optimal ventilator support and the correct interface to optimise ventilation and minimise side effects.<sup>3–5</sup>

## Key messages

- ▶ Ventilator data downloads facilitate optimisation of delivery of domiciliary NIV.
- ▶ Ventilator data downloads can facilitate early, objective assessment of ventilator prescriptions and leak.
- ▶ Ventilator data downloads can facilitate outpatient setups and are a useful interventional adjunct in the clinicians' toolbox.

In keeping with healthcare policies of providing high quality and cost-effective care in the community,<sup>6</sup> both numbers of people receiving and services delivering domiciliary NIV are increasing.<sup>7–9</sup> Traditionally, people undergo an inpatient admission for initiation of domiciliary NIV. With changing service models, more people are now being commenced on domiciliary NIV systems in outpatient or community settings. This has been shown to be as clinically effective as inpatient set-up in the neuromuscular disease (NMD) population<sup>10</sup>; cost effectiveness within the obese population is currently being assessed.<sup>11</sup> Improved patient experience, alongside reduced costs are other proposed benefits of outpatient initialisation of domiciliary NIV.<sup>11</sup> Nevertheless, a concern remains that domiciliary NIV outpatient set-up reduces the capacity for monitoring and assessing patients in comparison with bedside monitoring and the use of traditional tools such as transcutaneous CO<sub>2</sub> monitoring and arterial blood gases (ABGs).<sup>10</sup> Janssens *et al*<sup>12</sup> argue that ventilator data downloaded into specific software can be used as an alternative to support clinical decision-making in the successful management of domiciliary NIV. The use of ventilator data downloads



<sup>1</sup>Therapies Department, Royal Free London NHS Foundation Trust, London, UK

<sup>2</sup>Department of Thoracic Medicine, Royal Free London NHS Foundation Trust, London, UK

<sup>3</sup>Renaissance Research, Darlington, UK

<sup>4</sup>Department of Clinical Sciences, Brunel University London, London, UK

## Correspondence to

Stephanie K Mansell;  
stephanie.mansell1@nhs.net

may therefore help promote the wider use of outpatient set-up for home NIV; indeed ventilator download data, transmitted via a modem, can be described as a form of telemedicine.<sup>13</sup>

Data are available by both Secure Data (SD) card download and via modem technology, which means this data can be viewed remotely on a daily basis. Bench studies have demonstrated ventilator data parameters provided in manufacturers' software are reliable<sup>14 15</sup> and clinical studies have demonstrated the apnoea hypnoea index recorded by built in software is also reliable.<sup>16</sup> Most recently, Borel *et al*<sup>17</sup> demonstrated that ventilator download data, particularly respiratory rate, trigger % and usage, can predict an exacerbation of chronic obstructive pulmonary disease (COPD).

To date, there is minimal evidence for using data downloads in the assessment, treatment and management of people receiving domiciliary NIV.<sup>12 17–19</sup> The European Respiratory Society task force has called for further research in the area.<sup>13</sup> Therefore, this study aimed to assess the impact of using ventilator data downloads in the management of people with chronic hypercapnic ventilatory failure receiving domiciliary NIV.

Specifically, this study assessed the impact of ventilator data downloads on:

- ▶ Tidal volumes ( $V_T$ );
- ▶ The incidence of large leaks and interface issues;
- ▶ Individual compliance with domiciliary NIV;
- ▶ The incidence of continued hypercapnia (partial pressure of carbon dioxide ( $\text{PaCO}_2$ ) >6 kPa).

## METHODS

### Ethics

After discussion with the research and development department, it was deemed that ethical approval was not required for this study as all data were anonymous and collected as part of routine practice. Participants completed a data protection consent form for the installation of a modem. The study was registered with the trust's audit and governance department.

### Study design and participants

A longitudinal within-group repeated measures study design was used. All adult patients with chronic hypercapnic ventilatory failure receiving care under a domiciliary NIV service at the Royal Free London National Health Service (NHS) Foundation Trust were eligible for study inclusion. All patients were offered a review appointment between December 2014 and April 2015 and followed up by June 2015. Participants were excluded when they died during the study period, had their care transferred to another provider or declined to attend a review appointment.

### Procedure

Prior to the review appointment, all patients had been managed with clinical assessment, including symptoms and  $\text{PaCO}_2$  measurements obtained via Capillary Blood Gas (CBG); data from the patients' ventilators had not been used as part of their clinical management. The purpose of the review appointment was to:

- ▶ Exchange the participants' current home ventilator for a new model which was fitted with a modem.
- ▶ Use the ventilator data downloads to ensure optimal ventilation had been achieved, aiming for  $V_T$  of 8–10 mL/kg of ideal body weight.
- ▶ Review participants' interface via the leak measurement on the ventilator data download to ensure minimisation of both leak and side effects. Interfaces were changed or resized where participants had a leak of >60 L/min (as per manufacturer's guidance).
- ▶ Ensure participants and carers had received verbal and written patient information leaflets regarding their condition, use, care of and maintenance of their ventilator, interface and accessories.


Participants with  $V_T$  <6 mL/kg of ideal body weight or >10 mL/kg of ideal body weight were optimised through the use of a Philips Respironics A40 ventilator (Philips Respironics, Murrysville, USA) utilising Average Volume Assured Pressure Support/Average Volume Assured Pressure Support-Auto Expiratory Positive Airway Pressure (AVAPS/AVAPS-AE) mode and employing a titration approach.<sup>11</sup> In this study, the titration approach involved using AVAPS-AE mode over 4–6 weeks to evaluate the NIV prescription required by each participant.

All participants were monitored via the modem technology and EncoreAnywhere (Philips Respironics) system to measure and ensure:

- ▶ optimal ventilation, aiming for 8–10 mL/kg of ideal body weight;
- ▶ minimal leak;
- ▶ minimal side effects;
- ▶ optimal compliance, aiming for >4 hours/night for 70% of nights.

Data from the modems were reviewed by a consultant physiotherapist routinely twice weekly and when patients contacted the service. Participants were invited to a follow-up appointment on average 6 weeks post initial review appointment depending on appointment and patient availability. The purpose of this visit was to:

- ▶ Provide an appropriate ventilator for participants who required retitration: Philips Respironics A40, A30 or BiPAP ST (Philips Respironics).
- ▶ Ensure optimal ventilation had been achieved: aiming for  $V_T$  of 8–10 mL/kg of ideal body weight.
- ▶ Ensure interface was optimised to minimise both leak and side effects.
- ▶ Review patient compliance.



**Figure 1** Flow diagram of patient progress through the study. ITU, intensive treatment unit; NIV, non-invasive ventilation.

### Outcome measures

Ventilator data and PaCO<sub>2</sub> were collected by clinicians with appropriate skills and knowledge in domiciliary NIV but who had not provided care for the study participants.

### NIV data

Data were collected at the initial review appointment to act as the historical control for the cohort and further at the follow-up appointment 3–7 weeks later. Ventilator parameter data were collected via SD cards and subsequently modems to measure the following study outcomes: ventilator prescription, leak, V<sub>T</sub> (average over last 7 days of the data collection period) and compliance assessed as percentage of days NIV used >4 hours in the previous 28 days and average hours/day of NIV on days NIV was used.

### Capillary blood gas

PaCO<sub>2</sub> were obtained by trained clinicians via CBGs. CBGs were analysed on a RAPIDPoint 500 blood gas system (Siemens, UK). Where CBGs had been undertaken as part of the participant's normal care within 1 month, and

there had been no clinical change in the participant's status, CBG data were collected from clinic letters.

### Data analysis

Data were analysed using SPSS V.22 by an independent statistician. Descriptive statistics were used to report the demographics of the cohort. Data were analysed as paired data using a *t*-test. V<sub>T</sub>, respiratory rate, Ti and PaCO<sub>2</sub> levels were analysed using the paired *t*-test. Compliance, leak and ventilator prescription data were analysed with the Wilcoxon rank sum. The McNemar test was used to determine if there were differences in hypercapnic respiratory failure (defined as PaCO<sub>2</sub> >6 kPa). To assess if persistent hypercapnia was associated with leak, compliance or V<sub>T</sub>, an analysis of variance between participants who remained hypercapnic and participants where hypercapnia had resolved after review was undertaken.

## RESULTS

### Participants

At the start of the study, there were 78 participants. Data for 52 participants were analysed; data for the remaining

**Table 1** Type of ventilator before and after review appointment

Ventilator (Philips Respironics)	Before, % (n)	After, % (n)
A30	6% (3)	10% (5)
A40	–	33% (17)
BiPAP ST 30	19% (10)	58% (30)
Harmony	67% (35)	–
Synchrony	8% (4)	–
Interface		
Divers	2% (1)	8% (4)
Full face	27% (14)	31% (16)
Nasal	63% (33)	56% (29)
Pillows	8% (4)	6% (3)

15 were not available (see figure 1). Patients were excluded if they declined to attend a review appointment, as ventilator data for comparison were not available for these patients. Where patients were excluded due to mortality, they had not attended a review appointment nor were issued with a replacement ventilator.

There were more females in the group (men:women, 21:31) and the mean (SD) age of participants was 62 (18.8) years. The main reason for NIV use was obesity-related respiratory failure (33%), followed by COPD (27%), multiple diagnoses (13%), NMD (12%), chest wall deformity (10%) and other (6%). The median duration participants had been receiving domiciliary NIV prior to their review appointment was 33 months (IQR 15–48 months). The median time between the participants' review appointment and their last appointment in a respiratory clinic was 5.5 (IQR 1.6–9.7) months. At the first review

appointment, the mean nightly use of NIV was 6.53 hours and mean compliance was 90%, where compliance was defined as >4 hours/night NIV use for >70% of nights.<sup>20</sup> Prior to review, most participants used nasal (63%) and full face (27%) interfaces; post-review this was 56% and 31%, respectively (table 1).

Seventy-nine per cent (n=41) of patients required a change in their interface prescription, either in mask type or size. Eighty-seven per cent (n=45) of participants required an amendment to their ventilator prescription (table 2).

Prior to changing their prescription, seven individuals were achieving a  $V_T < 6$  mL/kg/ideal body weight and thus required an increase in the target  $V_T$ . Twenty-three individuals, although having a  $V_T > 10$  mL/kg/ideal body weight were thought to have been overventilated, and thus required titration of their ventilator prescription. Mean PaCO<sub>2</sub> at the start of the study was 6.59 (1.04) and post intervention was 6.45 (0.72) (mean difference 0.13, 95% CI -0.24 to 0.51, p=0.473).

#### $V_T$ and minute ventilation


$V_T$  measurements were available for 49 participants; the remaining participants were non-compliant and thus  $V_T$  measurements were not available. Results demonstrated a significant change in both actual  $V_T$  (mL) (mean difference 79.7 mL; 95% CI 12.1 to 147.2, p=0.022) and ideal  $V_T$  (mL/kg/ideal body weight) (mean difference 1.23; 95% CI 0.21 to 2.25, p=0.01) after prescription review (figure 2). There were two outliers whose  $V_T$  had been kept intentionally higher than usually expected for clinical reasons. When a reduction in  $V_T$  was trialled in these patients, they deteriorated and therefore their original prescription of >10 mL/kg ideal body weight was maintained. Following exclusion of the outliers,

**Table 2** Domiciliary non-invasive ventilation prescription pre-review and post-review

Prescription		Before, % (n) or median (IQR)	After, % (n) or median (IQR)	p value
Mode	ST	98% (51)	71% (37)	0.791
	S	2% (1)	–	
	AVAPS	–	8% (4)	
	AVAPS-AE	–	21% (11)	
IPAP, cm H <sub>2</sub> O for the whole data set		20 (15–24)	20 (18–24)	0.005
IPAP, cm H <sub>2</sub> O (Clinician set) n=35			20 (16–21)	
IPAP, cm H <sub>2</sub> O (AVAPS) n=18			24.5 (21.1–29.2)	
EPAP, cm H <sub>2</sub> O		4 (4–6)	6 (4–6)	<0.001
Pressure support, cm H <sub>2</sub> O		15 (10–18)	15.57 (12–18)	0.406
BPM		11 (11–18)	12 (12–12)	<0.001
Ti (s)		1.4 (1–1.5)	1.5 (1.5–1.8)	<0.001
Rise (100 ms)		1 (1–2)	3 (2–4)	<0.001

Data are presented as median (IQR) unless otherwise stated.

AVAPS, Average Volume Assured Pressure Support; BPM, breaths per minute; cm H<sub>2</sub>O, centimetres of water; EPAP, Expiratory Positive Airway Pressure; IPAP, Inspiratory Positive Airway Pressure; Ti, inspiratory time.



**Figure 2** Changes in tidal volume ( $V_T$ , mL/kg/ideal body weight) following non-invasive ventilation prescription review.

there remained a statistically significant change in both  $V_T$  (mL) (mean difference 91.1; 95% CI 27.4 to 154.8,  $p=0.006$ ) and ideal  $V_T$  (mean difference 1.38; 95% CI 0.42 to 2.34,  $p=0.006$ ). As demonstrated in [figure 2](#), there was a reduction in the range of the  $V_T$ . Where patients had a  $V_T$  of  $>10$  mL/kg/ideal body weight, there was a trend towards a reduction in their  $V_T$  (mean pre-review: 13.4 mL/kg/ideal body weight vs mean post-review: 9.7 mL/kg/ideal body weight). Where patients had a  $V_T$  of  $<6$  mL/kg/ideal body weight, there was a trend towards an increase in their  $V_T$  (mean pre-review 4.3 mL/kg/ideal body weight vs mean post-review 6.5 mL/kg/ideal body weight). At the initial review appointment, 15% of patients had a  $V_T$  of 8–10 mL/kg/ideal body weight versus 29% at the 6-week follow-up.

Minute ventilation measurements were available for 49 participants; the remaining participants were non-compliant and so minute volume measurements were not available. Results demonstrated an increase in minute ventilation after prescription review (mean difference 0.11 L/min; 95% CI  $-0.69$  to 0.92,  $p=0.781$ ).

### Leak


Leak measurements were available for 49 participants, where participants were non-compliant, leak measurements were not available. There was no change in leak following review of NIV prescriptions (mean (SD): 43 (23.4) L/min vs 45 (19.9) L/min,  $P=0.272$ ). Manufacturer's guidelines state that the ventilators used in this study are able to compensate for a leak of up to 60 L/min. Pre-review 12% ( $n=6$ ) of participants had a measured leak  $>60$  L/min, post-review only 4% ( $n=2$ ) continued to have a measured leak  $>60$  L/min.

### Compliance

Data were available for 50 participants. Compliance improved following a change in the NIV prescription; there was a significant increase in the percentage of days during which NIV was used for more than 4 hours per night (the recommended minimal use<sup>20</sup>, pre-prescription change: 90% vs post-prescription change: 96%,  $p=0.007$ ) ([figure 3](#)). There was an increase in the total number of hours used each night; however, this was not statistically significant (pre-prescription change: 6.53 vs post-prescription change: 6.94 hours,  $p=0.211$ ) ([figure 4](#)). Using a definition of compliance of  $>4$  hours use a day for 70% of days, prior to NIV prescription review 58% of participants were compliant and after review this increased to 70% ( $p=0.180$ ). Pre intervention, 19% ( $n=10$ ) of participants used their NIV for  $<4$  hours/night; in this subgroup, the mean use was 122 min/night, which increased to 270 min/night (mean difference 148 min; 95% CI 14 to 281,  $p=0.034$ ) post intervention for these people.

### Hypercapnic ventilatory failure

$\text{PaCO}_2$  data were available for only 21 participants (40%), as pre-review and post-review CBGs were only obtained if clinically indicated. There was a non-significant reduction in the incidence of ongoing hypercapnia (defined by a  $\text{PaCO}_2 >6$  kPa, pre-prescription change: 31% vs post-prescription change: 11%,  $p=0.125$ ) and measured  $\text{PaCO}_2$  (mean difference 0.14; 95% CI  $-0.24$  to 0.51,  $p=0.473$ ). A multivariate analysis showed no significant difference in leak ( $p=0.75$ ), compliance ( $p=0.65$ ) or  $V_T$  ( $p=0.65$ ) between participants who had persistent hypercapnia and those who were eucapnic. Individual inspection of the six participants with persistent hypercapnia showed they were either non-compliant ( $n=4$ , with average daily NIV use of 129 min) or had ongoing interface issues ( $n=2$ ).




**Figure 3** Compliance with non-invasive ventilation percentage of days with >4 hours use.

## DISCUSSION

These results demonstrate that the use of data downloaded via modems from domiciliary ventilators can lead to safe and clinically meaningful changes in domiciliary NIV prescription and patient outcomes. The increase seen in  $V_T$  is likely to be related to observed amendments in NIV prescriptions. The paired data show that there was a trend towards an increased IPAP and EPAP with a reduced rise time. There was also a change in the mean  $V_T$  post-review of 8.8 mL/kg/ideal body weight compared with a pre-review of 10.0 mL/

kg/ideal body weight; this may have been due to an improvement in leak in some individuals resulting in the delivery of more accurate  $V_T$ . Previous studies have reported  $V_T$  as mL/kg rather than mL/kg/ideal body weight, making comparison difficult. A target  $V_T$  of 8–10 mL/kg/ideal body weight has been employed in other studies; therefore, the  $V_T$  achieved in this study is comparable with other published studies.<sup>21</sup>

While not statistically significant, there was a reduction in the number of patients who had a leak greater than the manufacturer's set threshold (reduction in



**Figure 4** Compliance with non-invasive ventilation average hours/day before and after non-invasive ventilation prescription review.

66% of patients who had a leak greater than 60 L/min) and an observed reduction in the measured leak; this may have been due to changes in the interface prescription; however, the authors acknowledge that it is not possible to definitively attribute the leak reduction to this. In addition, since excessive leak may be identified by clinical features, such as dry mouth or excessive noise, the impact of the ventilator download data is difficult to judge in the absence of a control group. It is essential that leak is minimised to optimise ventilation; where leak is excessive (>60 L/min)  $V_T$  and efficiency of ventilation will be reduced.<sup>15</sup> Additionally, excessive leak can impact on patient comfort which may reduce compliance.<sup>22</sup> Furthermore, excessive leak can lead to overtightening of interface straps, which can cause pressure ulcers, a well-known side effect of NIV.<sup>22</sup> The patient-centred benefits of reduced mask leak could include reduced side effects such as eye irritation, reduced sleep disturbance from leak noise and potential reduced impact on bed partners sleep comfort. Thus, although a statistically significant reduction in the leak value was not seen, there may have been patient-centred gains, although the authors acknowledge that we cannot specifically infer this from the data collected. Future work specifically investigating the patient-centred benefits in terms of patient comfort, side effects and sleep quality from ventilator download-led prescription changes will help determine if the reduced leak has any clinical significance. Patient re-education on cleaning, caring and replacement of interfaces may have also contributed to the improvement in mask leak. Variation in algorithms used by different ventilator companies makes accurate comparison of interface leak across studies difficult.

Hours of NIV use was similar to that of other studies<sup>18</sup> and was greater than the 4 hours often suggested by clinicians. Although not statistically significant, a change in actual hours of daily NIV use was seen, along with an increase in participants' compliance, with a greater percentage of participants using NIV for more than 4 hours/night (90% vs 96%). In those patients who were considered to have low compliance, defined by less than 4 hours use of NIV on 70% of nights, there was an increase in the number of hours used on average by over 2 hours/night following download of ventilator data to adjust NIV prescription to ensure adequate ventilation, minimisation of leak and increase patient comfort. Suggesting that downloading ventilator data to assist in modifying NIV prescriptions is a useful adjunct to improve patient comfort and achieve a clinically meaningful increase in NIV compliance. Data from Masa *et al*<sup>23</sup> and Borel *et al*<sup>24</sup> suggest greater gains the longer NIV is used; thus a 2-hour increase in NIV use could result in greater patient benefit. Furthermore, the small sample size and heterogeneity of the population in this study may have resulted in a statistical underestimation of the changes observed in NIV compliance.

Overall, there was a reduction in use of pillows and nasal masks and an increase in prescription of full face and diver masks. Which suggests some patients may have been experiencing a significant leak (particularly from the mouth) or comfort issues which could have adversely affected compliance. It is likely that there was no one reason for increase in compliance and the cause is probably specific to each patient. Collecting patient comfort scores in future studies would provide more insight into this important area.

Measurements for PaCO<sub>2</sub> levels pre-review and post-review were available for 21 participants. While not statistically significant, there was an observed reduction in PaCO<sub>2</sub> and an overall reduction in the percentage of patients with ongoing hypercapnia, with 57% of patients having a reduction of their PaCO<sub>2</sub> to less than 6.5 kPa. We acknowledge that these data have been measured in a small number of patients and acknowledge that any conclusion on the effect of ventilator data downloads on PaCO<sub>2</sub> requires robust, prospective data.

### Limitations, bias and confounding factors

As this was a pragmatic clinically based study, there are a number of areas to be considered about the findings. First, the study design did not include a control group and thus the effect of the intervention is uncertain. Second, it was not feasible to blind the investigators collecting data, which may have resulted in observer bias; however, the objective nature of the data helps mitigate this risk.<sup>25</sup> To further minimise observer bias, investigators collecting data were not involved in participants' reviews. Third, this was a relatively small sample size in a heterogeneous population. Greater changes may have been seen in subgroups according to underlying disease pathology. Due to the novel nature of utilising ventilator data downloads, a sample size could not be calculated due to the paucity of previous literature. Additionally, the reliability of the software (EncoreAnywhere, Philips Respironics) has not been tested in bench studies, and reliability has not been reported. Therefore, the potential for measurement error is acknowledged. In addition, this software is only able to provide trend data for the measurement reported. Finally, we acknowledge the potential for the Hawthorne effect to have occurred as we were monitoring measurements not previously monitored.

Unrelated to the study, 12 participants died before their review appointment. Although attrition due to death is arguably inevitable in a study including domiciliary patients with NIV, it may nonetheless have contributed to study bias.

It was not possible to obtain CBG results pre-review and post-review for all participants; PaCO<sub>2</sub> data are therefore at risk of measurement bias and should be interpreted with caution. An independent and blinded statistician completed the data analysis, thus minimising reporting and observer bias. Multiple interventions were conducted at the review appointment, making it difficult to ascertain which

intervention had the biggest impact on compliance. Participants signed data protection consent forms for the application of a modem. They were therefore all aware that their NIV usage was being monitored. For some patients knowing that they were being monitored could cause behaviour change, thus, increasing compliance.

This study was designed to assess the clinical implications of utilising ventilator data downloads. The authors acknowledge that the lack of cost-effectiveness data is a limitation, however, a different study design would have been required to assess cost-effectiveness. For this study, assessment of the ventilator data downloads outside of the clinical appointments was undertaken by a consultant physiotherapist, the EncoreAnywhere system (Philips Respironics) has a built in flagging system, allowing clinicians' to prioritise patients with clinical concerns. Where a patient required a review of their ventilator data download, this was estimated to take approximately 3 min per patient. This may be offset by a reduced need for face-to-face clinical assessment. Future studies in this area should be designed to assess value in terms of cost as well as clinical effectiveness.

## CONCLUSION

Results from this study have demonstrated employing ventilator download functions can inform clinicians' decision-making to enable the optimal delivery of domiciliary NIV. Using ventilator data downloads in this outpatient setting facilitated an early, objective assessment of leak and ventilator prescriptions. Where outpatient NIV establishment and review are undertaken in lieu of traditional and more expensive inpatient environments, ventilator data downloads can provide clinicians with essential information, and therefore can be a useful interventional adjunct in the clinicians' toolbox.

**Twitter** @skmansell

**Acknowledgements** The authors wish to thank Robert Chase for his support during the study.

**Contributors** SKM and DDC conceived the idea for the project. SC and IH collected and inputted the data. KH and MW conducted the data analysis. SKM, SC, IH, MJW, KH, CK and SM were involved in the development of the manuscript.

**Funding** This project was partially funded by Philips Respironics. Funding was provided for data entry and statistical analysis. Third party statisticians (MW and KH) conducted the statistical analysis. Philips Respironics were not involved in the study design.

**Competing interests** SKM and SC have received sponsorship from Philips Respironics for conference attendance. SKM has provided consultation and speaker services to Philips Respironics.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** No additional data are available.

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

## REFERENCES

1. NICE. Motor neurone disease: the use of non-invasive ventilation in the management of motor neurone disease. (NICE) *NiHACE*. London, 2010.
2. Simonds AK, Elliott MW. Outcome of domiciliary nasal intermittent positive pressure ventilation in restrictive and obstructive disorders. *Thorax* 1995;50:604–9.
3. Bach JR, Robert D, Leger P, et al. Sleep fragmentation in kyphoscoliotic individuals with alveolar hypoventilation treated by NIPPV. *Chest* 1995;107:1552–8.
4. Meyer TJ, Pressman MR, Benditt J, et al. Air leaking through the mouth during nocturnal nasal ventilation: effect on sleep quality. *Sleep* 1997;20:561–9.
5. Teschler H, Stampa J, Ragette R, et al. Effect of mouth leak on effectiveness of nasal bilevel ventilatory assistance and sleep architecture. *Eur Respir J* 1999;14:1251–7.
6. NHS England. *Five Year Forward View*. London: NHS England, 2014.
7. Fletcher SV, Ewles S, Wilkinson JE. P226 The changing face of home NIV (non invasive ventilation). *Thorax* 2012;67(Suppl 2):A163.3–A164.
8. Mandal S, Suh E, Davies M, et al. Provision of home mechanical ventilation and sleep services for England survey. *Thorax* 2013;68:880–1.
9. Lloyd-Owen SJ, Donaldson GC, Ambrosino N, et al. Patterns of home mechanical ventilation use in Europe: results from the Eurovent survey. *Eur Respir J* 2005;25:1025–31.
10. Chatwin M, Nickol AH, Morrell MJ, et al. Randomised trial of inpatient versus outpatient initiation of home mechanical ventilation in patients with nocturnal hypoventilation. *Respir Med* 2008;102:1528–35.
11. Mandal S, Arbane G, Murphy P, et al. Medium-term cost-effectiveness of an automated non-invasive ventilation outpatient set-up versus a standard fixed level non-invasive ventilation inpatient set-up in obese patients with chronic respiratory failure: a protocol description. *BMJ Open* 2015;5:e007082.
12. Janssens JP, Borel JC, Pépin JL. Nocturnal monitoring of home non-invasive ventilation: the contribution of simple tools such as pulse oximetry, capnography, built-in ventilator software and autonomic markers of sleep fragmentation. *Thorax* 2011;66:438–45.
13. Ambrosino N, Vitacca M, Dreher M, et al. Tele-monitoring of ventilator-dependent patients: a European Respiratory Society Statement. *Eur Respir J* 2016;48:648–63.
14. Contal O, Vignaux L, Combescurie C, et al. Monitoring of noninvasive ventilation by built-in software of home bilevel ventilators: a bench study. *Chest* 2012;141:469–76.
15. Rabec C, Georges M, Kabeya NK, et al. Evaluating noninvasive ventilation using a monitoring system coupled to a ventilator: a bench-to-bedside study. *Eur Respir J* 2009;34:902–13.
16. Fernandez Alvarez R, Rabec C, Rubinos Cuadrado G, et al. Monitoring noninvasive ventilation in patients with obesity hypoventilation syndrome: comparison between ventilator built-in software and respiratory polygraphy. *Respiration* 2017;93:162–9.
17. Borel J-C, Pelletier J, Taleux N, et al. Parameters recorded by software of non-invasive ventilators predict COPD exacerbation: a proof-of-concept study. *Thorax* 2015.
18. Farr P, Bourqui P, Janssens P-P. What does built-in software of home ventilators tell us? *An observational study of 150 patients on home ventilation* 2011.
19. Cascón J, Fernández R, Quezada A, et al. Impact of built-in software analysis on monitoring patients under home mechanical ventilation. *Eur Resp J* 2013;42(Suppl 57):P3856.
20. Weaver TE, Maislin G, Dinges DF, et al. Relationship between hours of CPAP use and achieving normal levels of sleepiness and daily functioning. *Sleep* 2007;30:711–9.
21. Murphy PB, Davidson C, Hind MD, et al. Volume targeted versus pressure support non-invasive ventilation in patients with super obesity and chronic respiratory failure: a randomised controlled trial. *Thorax* 2012;67:727–34.
22. Elliott MW. The interface: crucial for successful noninvasive ventilation. *Eur Respir J* 2004;23:7–8.
23. Masa JF, Corral-Penafiel J. Should use of 4 hours continuous positive airway pressure per night be considered acceptable compliance? *Eur Respir J* 2014;44:1119–20.
24. Borel JC, Pepin JL, Pison C, et al. Long-term adherence with non-invasive ventilation improves prognosis in obese COPD patients. *Respirology* 2014;6:857–65.
25. Deeks JJ, Higgins J, Altman DG. *Analysing data and undertaking meta-analyses. Cochrane handbook for systematic reviews of interventions: Cochrane book series*, 2008:243–96.