Is childhood wheeze and asthma in Latin America associated with poor hygiene and infection? A systematic review

Cristina Ardura-Garcia,¹ Paul Garner,¹ Philip J Cooper^{2,3}

To cite: Ardura-Garcia C, Garner P, Cooper PJ. Is childhood wheeze and asthma in Latin America associated with poor hygiene and infection? A systematic review. *BMJ Open Res* 2018;**5**:e000249. doi:10.1136/bmjresp-2017-000249

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjresp-2017-000249>).

Received 8 November 2017
Revised 26 January 2018

¹Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, UK

²Facultad de Ciencias Medicas, de la Salud y la Vida, Universidad Internacional del Ecuador, Quito, Ecuador

³Institute of Infection and Immunity, St George's, University of London, London, UK

Correspondence to

Dr Cristina Ardura-Garcia, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, L3 5QA, UK; crisardura@gmail.com

ABSTRACT

Introduction High asthma prevalence in Latin-American cities is thought to be caused by poor hygiene and infections. This contradicts the widely accepted 'hygiene hypothesis' for asthma aetiology.

Methods Systematic review of observational studies evaluating the association between poor hygiene exposures or infections and asthma/wheeze among Latin-American children aged 4–16 years. MEDLINE, EMBASE, LILACS and CINAHL electronic databases were searched following a predefined strategy to 18 December 2017. We quantified outcomes measured and reported, assessed risk of bias and tabulated the results.

Results Forty-five studies included: 6 cohort, 30 cross-sectional and 9 case-control studies. 26 cross-sectional studies were school-based surveys (14 of over 3000 children), whereas 5 case-control studies were hospital/health centre-based. Exposures measured and reported varied substantially between studies, and current wheeze was the most common outcome reported. Data showed selective reporting based on statistical significance (P value <0.05): 17/45 studies did not clearly describe the number of exposures measured and 15/45 studies reported on less than 50% of the exposures measured. Most exposures studied did not show an association with wheeze or asthma, except for a generally increased risk associated with acute respiratory infections in early life. Contradictory associations were observed frequently between different studies.

Conclusion Selective reporting is common in observational studies exploring the association between environmental exposures and risk of wheeze/asthma. This, together with the use of different study outcomes (wheeze/asthma) associated with possibly distinct causal mechanisms, complicates inferences about the role of poor hygiene exposures and childhood infections in explaining asthma prevalence in Latin-American children.

INTRODUCTION

Asthma prevalence has increased worldwide¹ and is estimated to affect 400 million people.² A widely accepted explanation for increased asthma prevalence in industrialised countries is provided by the 'hygiene hypothesis' in which diminished exposures to certain

micro-organisms during the early years of life are purported to increase allergic disease risk.³

The immunological mechanisms underlying the hygiene hypothesis remain under debate. There is evidence that micro-organisms such as those present in the gut microbiota, nasal colonisers or intestinal helminths drive regulatory immune cells to maintain immune homeostasis. Consequently, reduced or altered exposures to such micro-organisms may lead to a failure in immune regulation, thus increasing the risk of chronic inflammation.⁴

Asthma is a heterogeneous disease caused by complex gene-environment interactions. It appears reasonable to attribute increasing asthma prevalence—occurring over a relative period of time—to changes in environmental exposures rather than in gene frequencies in human populations. Several studies have demonstrated a reduced risk of atopy and allergic diseases associated with farming,⁵ helminth infestations⁶ and contacts with other children. However, findings between studies are not always consistent,⁷ perhaps explained by the use of different asthma definitions, phenotypes (eg, atopic vs non-atopic) and diverse comparison groups.⁸ Inconsistent findings between studies may reflect also different underlying mechanisms and associated environmental exposures. Asthma is a complex disease likely consisting of several phenotypes, of which the most widely used are defined by the presence and absence of atopy. The proportion of asthma cases attributable to atopy is positively associated with economic development,⁸ and research in industrialised countries has mainly focused on atopic asthma, whereas non-atopic asthma, the predominant childhood asthma phenotype in Latin America,^{8–10} remains understudied.

Asthma prevalence may now be increasing in formerly low-risk, low-income and middle-income countries, while it has reached a plateau in many high-risk, high-income countries.¹¹ High rates of asthma have been reported in some Latin-American cities¹¹ in which conditions of overcrowding, poor hygiene and high infectious disease burdens predominate. Further, higher rates of asthma have been described in poorer Latin-American regions with a high prevalence of acute respiratory infections and intestinal parasite infestation in early childhood.¹² Several Latin-American studies have shown that factors associated with poor hygiene may be associated with a higher risk of non-atopic asthma or wheezing.^{9 13 14} The role of chronic infections (eg, intestinal parasites) in the development of asthma in Latin America remains controversial: such exposures attenuate atopy but appear to have little impact on asthma prevalence.^{15 16}

Our aim was to summarise and appraise the evidence of association between asthma or wheeze and (1) poor hygiene and (2) past and current parasite infections and chronic viral or bacterial infections.

METHODS

Inclusion criteria

Studies were included if they met the following criteria (Annex 1): (1) observational study design (cross-sectional, cohort and case-control studies); (2) children aged 4–16 years, born and currently living in a Latin-American country; (3) asthma (guidelines criteria or reported doctor's diagnosis), wheeze reported by written or video questionnaire ('has your child/have you had wheezing during the last 12 months?') or a doctor's diagnosis, or bronchial hyper-responsiveness included as outcomes and (4) environmental exposures associated with a higher risk of infection or infections (gastrointestinal or respiratory infections, current/past intestinal parasites or chronic bacterial or viral infections) listed as exposures.

Exclusion criteria were: (1) reports not in English, Spanish, French or Portuguese; (2) published before 1980; (3) children outside Latin America involved as participants; (4) acute wheeze or asthma used as outcomes and (5) unpublished data and conference abstracts.

Data sources and searches

We identified relevant studies by searching MEDLINE, EMBASE, LILACS and CINAHL electronic databases. No language, time or publication status restrictions were applied, and the last search was run on 28 December 2017 by CA-G together with an information specialist from Cochrane Infectious Diseases Group (online supplementary material: Annex 1). We used reference manager software (EndNote) to merge all the search results and remove duplicates.

Titles and abstracts were screened by CA-G to exclude studies. When abstracts were not available, tables and

most relevant content were reviewed. Finally, full-text articles of selected papers were reviewed for eligibility as described above.

Data extraction

We piloted our initial data extraction form with nine of the most relevant studies to develop the final form, which was used to retrieve data from selected studies. Data extraction was carried out by CA-G as prespecified in the study protocol and included: study characteristics (design, location, year and duration), participants (number, age range, sample selection and method of recruitment), outcomes (definitions and prevalence in study sample), exposures (type, methods used to assess exposure, time and duration of exposure and number of exposures measured) and results. Data were summarised in tables, and studies grouped by design. Where possible, results were presented and divided into atopic and non-atopic wheeze or asthma, with atopy defined as a positive skin prick test or positive specific serum IgE to any aeroallergen.

Risk of bias assessment and data analysis

Potential risk of bias was described in relation to (1) sampling (random sampling and response rates); (2) reporting of the results (number of exposures described and reported to evaluate selective reporting) and (3) statistical analysis (statistical corrections for multiple significance testing). We reported if studies were adjusted for potential confounders and effect modifiers.

Synthesis

We examined results by exposures related to the home environment, animal contact, contact with children, early-life infections and current/past infection with intestinal parasites. Results were reported together with the outcome used (wheeze or asthma).

RESULTS

The search yielded a total of 860 reports of which 125 full-text articles were assessed for eligibility. Sixty reports representing 45 studies fulfilled the eligibility criteria (figure 1).

Description of studies

Study design

The main characteristics of included studies are summarised in table 1. There were 6 cohort, 30 cross-sectional and 9 case-control studies. Twenty-seven (60%) studies were from Brazil, and the majority were urban-based, done between 1987 and 2014 (online supplementary table 1).

Of the cross-sectional studies, 14 had a sample size of greater than 3000 subjects as recommended by the International Study of Asthma and Allergies in

Figure 1 Flow diagram of included and excluded studies.

Childhood (ISAAC),¹⁷ whereas the rest ranged 100–2700 subjects. The majority were school-based.

As for the case–control studies, 6 were done in hospitals or health centres and 3 in schools or homes with samples varying: 400–600 children in 4 studies, 100–200 in 4 and 19 in 1.

Outcomes and case definition

The most frequent outcome was current or recent wheeze, as defined by the ISAAC question: ‘Has your child/have you had wheezing or whistling in the chest in the last 12 months?’.¹⁸ This was used in 5 cohort and 20 cross-sectional studies and was the case definition in 3 case–control studies. Some studies representing more than one report varied the outcome used (see footnote in online supplementary table 1).

Case definition for the remaining case–control studies was either doctor diagnosis of asthma or that defined by the Global Initiative for Asthma guidelines.¹⁹ Study

outcomes used are provided in online supplementary table 1.

Measurement of exposures

Table 1 shows the exposures measured. Exposures relating to a higher risk of infection or specific infections varied greatly between studies. Although 5 of the cohorts, 20 of the cross-sectional and 4 of the case–control studies used ISAAC-derived questionnaires, the exposures measured differed greatly. Social Changes, Asthma and Allergy in Latin America (SCAALA) Brazil^{9 15 16 20 21} collected stool and blood samples to diagnose intestinal parasites and viral or bacterial infections. Two more cohort,^{22–26} 10 cross-sectional and 4 case–control studies analysed intestinal parasites (3 using serology). One other cohort²⁷ and one cross-sectional study²⁸ also collected blood to detect viral or bacterial infections.

Only two cohort^{9 15 16 20–22 25} and two cross-sectional^{10 29–31} studies measured at least one exposure variable in the

Table 1 Classification of exposures related to poor hygiene and infection

	Definition	Total (45)	Cohort (6)	Cross-sectional (30)	Case-control (9)
Home environment	Water and sanitation, garbage disposal, house cleaning, housing construction, endotoxins	9	2	6	1
Animal contact	Contact with pets, farm animals and cockroaches inside/outside the house	34	5	23	6
Contact with other children	Overcrowding in home, day-care attendance and having older siblings	22	6	11	5
Early-life infection	Acute respiratory and gastrointestinal infections during infancy, worm infections	11	4	6	1
Demonstrated infection	Intestinal parasites or chronic viral or bacterial infections diagnosed by serology or stool sample	21	4	11	6

Numbers represent the number of studies that measured at least one exposure related to each category, even if the results were not published.

five exposure categories. The most frequently measured exposure category was animal contact followed by contact with other children and infections (table 1).

Risk of bias

Sampling

Sampling methods and response and follow-up rates are shown in online supplementary table 1. Ten of 30 cross-sectional studies used non-random (or unclear selection) samples. Eighteen of these 30 studies obtained a response rate $\geq 84\%$, 5 between 45% and 75% and 7 did not provide response rates.

Reporting of results

The risk factor questionnaire for the ISAAC Phase II study included 67 variables (10 hygiene or infection-related), whereas Phase III included 34 for 13–14 years (3 hygiene or infection-related) and 43 for 6–7 years (7 hygiene or infection-related).

One-third of papers (17/45) did not clearly describe number of exposures measured (table 2). Eleven (39%) of 28 studies that described measured variables reported on $<50\%$ of them, 3/28 (11%) reported on 50%–80% and 14/28 (50%) reported on more than 80% of the measured exposures. Among all studies: 15/45 (33%) reported on $<50\%$ of measured exposures, 8/45 (18%) on 50%–80%, 18/45 (44%) on $>80\%$ and 4/45 had no data.^{32–35}

The most common criteria used for presentation of results were statistical significance variables (P value <0.05 or <0.2) in univariate analyses which were then included in multivariate models (table 2).

Statistical analysis

None of the reports carried out any corrections for multiple significance testing.

Measurement of confounders and effect modifiers

Thirty-six per cent (16/45) of papers did not show results adjusted for potential confounders, or information on adjustment was unclear (table 2).

Adjustment for several risk factors for asthma identified in previous published literature,^{10 36–41} which could act as confounders or effect modifiers are represented in online supplementary table 2. None of the studies were adjusted for all of the possible confounders (ie, age, gender, atopy, bronchiolitis in infancy, parental asthma, breastfeeding, socioeconomic status and tobacco exposure). Age, gender and parental asthma were most frequently adjusted for.

Atopy and history of bronchiolitis may behave as effect modifiers when studying risk factors for asthma.^{10 38} SCAALA Brazil,^{9 15 16 20 21} ECUAVIDA,^{22 25} SCAALA Ecuador^{10 30 31} and the Uruguiana study^{13 14} represented results stratified by atopy, and another 13 studies adjusted for atopy. Only five studies adjusted for

bronchiolitis. Some reports did not distinguish bronchiolitis from early-life respiratory infections.

Results by environmental exposures

The high heterogeneity in methodology, exposures studied and outcomes measured precluded a meta-analysis. The main results are summarised in table 3. The results for the most relevant exposures are provided in online supplementary tables 3–7.

Home environment

Water and sanitation infrastructure, garbage disposal, frequency of house cleaning, housing construction materials and presence of endotoxins were analysed in nine studies (table 1). Most results showed no association, except for an increased risk of current wheeze with open-field defecation compared with toilets or latrines (adjusted OR (AOR) 1.31, 95% CI 1.02 to 1.68)¹⁰ and with lack of potable drinking water (AOR 1.44, 95% CI 1.16 to 1.78),³⁰ an increased risk of asthma with the presence of sewage disposal⁴² and increased risk of non-atopic wheeze compared with non-atopic non-wheezing with infrequent house cleaning.⁹

Animal contact

The association between animal contact and wheeze was investigated in 34 studies (table 1), though only reported in 30 studies (online supplementary table 3). All studied the effect of domestic animals (11 with cats and 9 with dogs), with inconsistent findings. Farm animals were evaluated in 10 studies, one showed a decreased risk of asthma after regular contact,⁴³ another showed an increased risk of cumulative asthma in boys aged 6–7 years after maternal contact with animals during pregnancy,⁴⁴ with no associations (or no results presented) in 8 other studies. Overall, there was no consistent effect that might reflect protection or causality.

Contact with other children (older siblings, overcrowding and day-care attendance)

Of 12 studies reporting the relationship with having older siblings, there was no obvious pattern (online supplementary table 4). The 13 studies evaluating the effect of overcrowding did not report on an association, except for Brandão cohort,⁴⁵ which showed an increased risk of current wheeze at 6 years among children born in private hospitals and not exposed to household overcrowding (AOR 6.46, 95% CI 1.11 to 37.57)⁴⁵ and no association at 13–14 years.⁴⁶ Day-care attendance increased the risk of non-atopic wheeze in a cohort study⁹ and of asthma in a case-control study,⁴⁷ decreased the risk of asthma in another case-control study⁴³ and showed no association with wheeze or asthma in eight other studies (online supplementary table 5).

Table 2 Risk of bias: reporting of exposures measured, adjustment for confounders and proportion of variables associated with asthma or wheeze

Design	Study	ISAAC questionnaire	Exposures measured	Exposures reported	Criteria for presentation	Adjustment for confounders	Hygiene exposures with association (P<0.005)	Non-hygiene exposures with association (P<0.05)	Total exposures with association (P<0.005)
Cohort	Brandão et al ^{45,46}	Yes	43 (6–7 years)? 34 (13–14 years)?	26 21	Not described for univariate analysis. Variables reported varied between the different ages	Yes	3/7	3/15	6/22
	SCAALA Brazil ^{9,15,16,20,21}	Yes	55	50	*Meaningfully associated* in univariate or multivariate analysis (with <i>Schistosoma mansoni</i> and hookworm excluded from analysis)	Yes	4/28	1/22	5/50
	Pelotas cohort ^{23,24,26}	Modified	20?	20	Not described for multivariate. No hygiene exposure in multivariate*	Yes*	1/5	7/15	8/20
	Cuban study ^{60–62}	Modified	25	24	One more variable only used as confounder in multivariate	Yes	2/15	6/10	8/25
	ECUAVIDA ^{22,25}	Modified	47?	25	Not described, except confounders chosen for significant associations with the 5-year AW phenotype and on previously reported associations with asthma or with microbiome shifts	Yes	3/12	4/13	7/25
Cross-sectional	Zepeda et al ²⁷	No	15	14	Not described	No	1/12	0/3	1/15
	SCAALA Ecuador ^{10,30,31}	Yes	47	27	P<0.2 in univariate analysis or included in multivariate analysis. Parasites with low prevalence not represented	Yes	5/15	2/32	7/47
	Uruguiana study ^{13,14}	Yes	41?	11–12	Risk factors known to be associated with wheeze/asthma and those significantly associated in the bivariate analysis. Intestinal parasites	Yes†	1/11	6/30	7/41?
	Azalim et al ³²	Modified	?	12	Not described	Yes	0/2	4/11	4/13
	Barraza et al ³³	Yes	?	16	Not described	Yes	2/3	8/13	10/16
	Barreto and Sole ²⁸	Yes	33	15	P<0.2 in univariate analysis and P<0.05 in bivariate analysis	No	3/?	12/?	15/33
	Bragagnoli and Silva ⁶³	Yes	18	15	Only presented results for the <i>Ascaris lumbricoides</i> coinfections with other parasites (not those other parasites alone)	No	3/14	2/4	5/18
	Casagrande et al ²⁹	Yes	33?	31	Not described (P<0.2 in univariate analysis to include in bivariate)	Yes	0/11	2/20	2/31
	Cooper et al ⁶⁴	Yes	4	3	Not specified (low proportion of <i>Ancylostoma duodenale</i> ?)	Yes	0/3	–	0/3

Continued

Table 2 Continued

Design	Study	ISAAC questionnaire	Exposures measured	Exposures reported	Criteria for presentation	Adjustment for confounders	Hygiene exposures with association (P<0.005)	Non-hygiene exposures with association (P<0.05)	Total exposures with association (P<0.005)
	Del-Rio-Navarro <i>et al</i> ⁴⁴	Yes	43 (6–7 years) 34 (13–14 years)	6–10 7–14	Statistically significant results (for univariate and multivariate)	Unclear	1/1 8/13	7/9 1/1	8/10 9/14
	Endara <i>et al</i> ⁶⁵	Yes	1 (+6 parasites)	1	No association with intestinal parasites	Yes	0/1 (7)	0	0/1 (7)
	Freitas <i>et al</i> ⁶⁶	Yes	65	10	P<0.2 in univariate was included in multivariate. Final model variables P<0.05	Yes	1/10	2/55	3/65
	Garcia <i>et al</i> ⁶⁷	Yes	35	12	P<0.25 in bivariate analysis	Yes	1/3	5/32	6/35
	Gomes de Luna <i>et al</i> ⁶⁸	Modified	34	34		Yes	0/1	4/33	4/34
	Guimarães <i>et al</i> ⁶⁹	No	14	14		No	0/2	2/12	2/14
	Hagel <i>et al</i> ⁷⁰	–	2	2		No	0/2	–	0/2
	Han <i>et al</i> ^{71 72}	Modified	37?	21/26	Not specified	Yes§	2/13	11/24	13/37
	Kuschnir and Alves da Cunha ⁷³	Yes	34	10	Not stated	Yes	2/3	3/31	5/34
	Lima <i>et al</i> ⁷⁴	Modified	9	9		Yes	0/1	3/8	3/9
	Maia <i>et al</i> ⁷⁵	No	8	3	P<0.05 in final multivariate model	Yes	1/1	2/7	3/8
	Palvo <i>et al</i> ⁷⁶	Modified	14	9	P<0.2 in univariate analysis and P<0.05 in multivariate analysis	Yes	1/4	5/10	6/14
	Prietsch <i>et al</i> ⁷⁷	No	28?	17	P<0.2 in univariate analysis (to be included in multivariate)	Yes	1/6	6/22	7/28
	Quiroz-Arcoteles <i>et al</i> ⁴²	No	15?	5	Not stated	No	4?/5	6?/10	10?/15
	Ribeiro <i>et al</i> ⁷⁸	No	17	7	Not stated	No	0/5	1/12	1/17
	Rojas Molina <i>et al</i> ³⁴	Modified	?	3–4	Only significant variables	No	1/?	2–3/?	3–4/?
	Silva <i>et al</i> ⁷⁹	No	1	1		Yes	0/1	–	0/1
	Solis-Soto <i>et al</i> ³⁵	Yes	?	9	Not specified	Yes	1/5	2/4	3/9
	Soto-Quiros <i>et al</i> ⁸⁰	No	10	10		No	1/1	4/9	5/10
	Souza <i>et al</i> ⁸¹	–	1	0	No result for asthma alone (only for respiratory allergy)	No	0/1	–	0/1
	Tintori <i>et al</i> ⁸²	No	14?	14	P<0.5 in univariate was included in multivariate. Final model variables P<0.05	Yes	1/2	11/12	12/14
Case-control	Toledo <i>et al</i> ⁸³	Modified	12	4	Not stated	No	0/4	0/8	0/12
	Boneberger <i>et al</i> ⁴³	Modified	13	11	Matched by sex and age (not represented)	Yes	3/6	1/7	4/13
	Cadore <i>et al</i> ⁸⁴	No	14?	14		Yes	1/4	5/10	6/14

Continued

Table 2 Continued

Design	Study	ISAAC questionnaire	Exposures measured	Exposures reported	Criteria for presentation	Adjustment for confounders	Hygiene exposures with association (P<0.005)	Non-hygiene exposures with association (P<0.05)	Total exposures with association (P<0.005)
	Coelho <i>et al</i> ⁴⁷	Yes	67?	35	Not stated. P<0.2 in bivariate included in multivariate	Yes	2/11	4/24	6/35
	Jucá <i>et al</i> ⁸⁵	Yes	67?	26	Not stated. P<0.2 in bivariate included in multivariate	Yes	1/10	7/58	8/68
	López <i>et al</i> ⁸⁶	–	1	1		No	0/1	–	0/1
	Mendoza <i>et al</i> ⁸⁷	–	6	6		No	1/6	–	1/6
	Moraes <i>et al</i> ⁸⁸	No	22	22		Unclear [¶]	1/3	1/19	2/22
	Oliveira-Santos <i>et al</i> ⁸⁹	Yes	67?	26	Not stated. P<0.25 in bivariate included in multivariate	Yes	2/6	1/20	3/26
	Rizzo <i>et al</i> ⁹⁰	No	3	0	'No significant difference' in text, but no numbers represented	No	0/2	0/1	0/3

? : unclear number or not included.

*Muñoz *et al*²⁶: no description of variables measured and represented. No adjustment.

†Only for Pereira *et al*¹⁴ not da Silva *et al*.¹³

‡Same questionnaire as SCAALA Ecuador.

§Only rural residence and antibiotic use and bronchiolitis in the first year of life were adjusted for in multivariable analysis, as was the objective of the study.

¶Not published.

Exposures associated to wheeze or asthma: shown over total exposures measured or represented (when number of exposures measured was not clear). When results presented were divided in subgroups, highest number of associated variables from any of the subgroups was selected for the table.

AW, atopic wheeze; ISAAC, International Study on Asthma and Allergies in Children; SCAALA, Social Changes, Asthma and Allergy in Latin America.

Table 3 Results for the association between exposures related to a higher risk of infection and wheeze/asthma

Study	Home environment				Animal contact			Contact with other children				Early-life infection		Demonstrated infection						
	House*	Sanitation†	Water‡	Endotoxin	Housing	Garbage§	Animal contact		Farm animal	Insects¶	Overcrowding	Day care	Older sibling	ARI	GI	Intestinal parasite				
							Inside	Outside								Pets	IgG	Virus**	Bacteria**	
Brandão																				
SCAALA Brazil	↑ ²	NA	NA			NA	↑ ²	NA	–	NA	NA/↑ ¹	NA	↑ ²	NA	NA	NA	NA ³	NA	NA	NA
Pelotas cohort							NA ⁴		NA ⁴	NA	NA	NA	↑ ⁵	NA ⁵						
Cuban study									NA	NA	NA	NA				NA/↓ ↑ ⁶				
ECUAVIDA		NA	NA					NA	NA	NA	NA	NA	↑	NA	NA	↓/↑ ⁷				
Zepeda et al ⁶⁷											NA	NA						NA		
SCAALA Ecuador	↑ ⁸	NA/↑ ⁹			NA ⁹		–	NA	NA/↑ ¹⁰	NA	NA	NA	NA ¹⁰ /↓ ¹¹			NA/↓ ↑ ¹²				
Uruguiana study							–	–	–	–	–	–	NA/↓ ¹³			NA/↓ ↑ ¹⁴				
Azalim et al ⁶²									NA		NA									
Barraza et al ⁶³								NA					↑							
Barreto and Solé ²⁸									↓											↑ ¹⁵
Bragagnoli and Silva ⁶³																NA/↓ ↑ ¹⁶				
Casagrande et al ²⁹			NA				–	–	NA	–	NA	NA	NA	NA	NA	NA				
Cooper et al ⁶⁴									↑	↑										
Del-Rio-Navarro et al ⁴⁴																				
Endara et al ⁶⁵									↑				↑							
Freitas et al ⁶⁶									↑		NA									
Garcia et al ⁶⁷									↑ ¹⁷	–										
Gomes et al ⁶⁸									NA											
Guimarães et al ⁶⁹											NA									
Hagel et al ⁷⁰								NA												
Han et al ⁷¹									NA	NA		NA	NA ¹⁸	↑						
Kuschnir and Alves da Cunha ⁷³									↑				↓							
Lima et al ⁷⁴									NA											
Maia et al ⁷⁵									↑											
Palvo et al ⁷⁶									↑			–	–							
Prietsch et al ⁷⁷		NA	–		–				–		NA			↑						
Quiroz-Arcenales et al ⁴²		↑ ¹⁹	NA			↑	–	–						↑						
Ribeiro et al ⁷⁸					–				NA		–	NA	–							
Rojas et al ⁶⁴								↑					–							
Silva et al ⁷⁹																				NA
Solis-Soto et al ⁶⁵		NA ²⁰				NA ²⁰			NA	NA	↑ ²¹									

Continued

Table 3 Continued

Study	Home environment				Animal contact			Contact with other children				Early-life infection		Demonstrated infection					
	House*	Sanitation†	Water‡	Endotoxin	Housing	Garbage§	Inside	Outside	Pets	Farm animal	Insects¶	Overcrowding	Day care	Older sibling	ARI	GI	Intestinal parasite		
																	Stool	IgG	Bacteria**
Soto-Quiros et al ⁶⁰															↑				
Souza et al ⁶¹									NA/↑ ²²									NA	
Tintori et al ⁶²																			
Toledo et al ⁶³							NA	NA			NA								
Boneberger et al ⁶³									NA	↓			↓	NA	↑				
Cadore et al ⁶⁴									NA/↑ ²³			NA						NA	
Coelho et al ⁶⁷							NA		NA			NA	NA/↑ ²⁴	NA				NA	
Jucá et al ⁶⁵							↑	-				-	-	-					
López et al ⁶⁶																		NA	
Mendoza et al ⁶⁷																		↓	
Moraes et al ⁶⁸									↓			NA						NA	
Oliveira-Santos et al ⁶⁹									NA/↑ ²⁵			NA	NA	↓				NA ²⁵	
Rizzo et al ⁶⁰											NA								

1: increased risk of current wheeze at 6 years; ⁴⁵ no association at 13–14 years of age when adjusted; ⁴⁶ 2: increased risk of non-atopic wheeze compared with non-atopic wheeze; ⁴⁷ animals inside; rats; 3: only *Toxocara* IgG increased risk of atopic wheeze when compared with non-atopic non-wheezers; ⁴⁸ 4: only in Chatkin *et al.*⁴⁵; 5: only in Muñoz *et al.*, 2008; 6: current helminth infection; ⁴⁹ past *Ascaris lumbricoides*, *Trichuris trichiura* and hookworm infection (Werf 2013) were NA, history of *Ascaris* infection increased the risk of current wheeze; ⁵⁰ 7: maternal geohelminths increased the risk of current wheeze, childhood geohelminths to 36 months decreased risk of current wheeze and asthma; 8: NA in Cooper *et al.*⁵⁰; 9: increased risk of wheeze with lack of potable drinking water in Cooper *et al.*⁵⁰; 10: dog inside the house increased risk of current wheeze in urban setting; 11: only in Cooper *et al.*⁵⁰; 12: no association for any geohelminth, hookworm or *Ascaris*^{10,31,30} decreased risk of atopic wheeze with *T. trichiura* infection^{10,31}; 13: decreased risk of active asthma, NA for wheeze; 14: high load of *Ascaris* increased risk of active asthma; ¹⁴ *Giardia* infection decreased risk and high load of helminth infection increased risk; ¹⁵ history of measles or tuberculosis increased the risk of asthma; 16: the risk of wheeze decreased with light *A. lumbricoides* infections and increased with heavy infections and with *T. trichiura* coinfections; 17: no association in 13–14 year group; 18: bronchiolitis increased risk of current wheeze and asthma in 6–7 years; 19: absence of sewage disposal increased the risk of asthma; 20: precarious household conditions together as one exposure including: precarious floor, precarious walls, precarious source of water, precarious sewage system; 21: presence of disease vectors at home: fleas, ticks, kissing bugs, mice, bedbugs, flies; 22: dogs in the house increased the risk of current wheeze, no association for cats; 23: contact with cats increased risk of asthma, with dogs had no association; 24: kindergarten increased risk, day care was not associated; 25: dogs inside the house currently not associated, during 1 year of life decreased the risk of current wheeze. Worm infection in the past not associated with current wheeze.

*Infrequent house cleaning.

†No toilet or latrine.

‡No clean drinking water.

§No garbage disposal.

¶Cockroaches in the house.

**Positive serology for *Helicobacter pylori*, hepatitis A virus, herpes zoster virus, herpes simplex virus, Epstein-Barr virus.

ARI, acute respiratory tract infection; GI, gastrointestinal; SCAALLA, Social Changes, Asthma and Allergy in Latin America.

–: exposure measure not represented; NA: no association ($P \leq 0.05$); ↑: increased risk of asthma (OR < 1, $P < 0.05$); ↓: decreased risk of asthma (OR < 1, $P < 0.05$).

Early-life infections

Online supplementary table 6 shows the effects of early-life infections on wheeze and asthma. Nine studies demonstrated an increased risk of wheeze and asthma (both atopic and non-atopic) associated with acute respiratory infections in early life, though the cohort study that reported AOR found an association with non-atopic wheeze and not with atopic wheeze.⁹ A further cross-sectional study found no association between viral bronchiolitis and recurrent wheezing.²⁷ Gastrointestinal infections in early life showed no association with wheeze or asthma in three studies.^{9 22 26}

Intestinal parasites

Intestinal parasites were analysed in 18 studies (online supplementary table 7), reporting no association with wheeze or asthma in 11. Positive and negative associations reported in the remaining seven studies varied greatly depending on the specific parasite, load of infestation, presence of coinfections and age of exposure (intrauterus, early life or current).

DISCUSSION

Overall, current evidence is not sufficient to derive a conclusion as to whether poor hygiene exposures and early-life infections affect the risk of developing childhood wheeze or asthma in Latin America. Only six cohort studies were included in this review, five of which followed up children from the first few years of life, though only one was specifically designed to study asthma outcomes. Selective reporting of statistically significant results was common to many studies (with the exception of the cohort studies), exposure variables measured varied greatly between studies and the majority of studies showed no associations with asthma or wheeze. The exception was early-life acute respiratory infections, which showed reasonably consistent positive associations with wheeze (mainly non-atopic) and asthma across studies.

The use of a wide literature search with no language restriction and including a Latin-American database probably identified the majority of relevant studies. The inclusion of studies from several South and Central American countries ensured the representation of different Latin-American regions. Most cross-sectional studies included in this review were methodologically of good quality following the ISAAC guidelines^{17 18} and included large sample sizes. The use of a widely validated questionnaire such as the ISAAC questionnaire^{18 48 49} in a large proportion of studies provided a reasonably standardised instrument to measure exposures and wheeze or asthma.

Substantial selective reporting was observed across studies, with a large or even unknown number of exposures studied and only statistically significant variables reported. Similarly, none of the studies applied any statistical correction for multiple testing, even though more than 30 variables were studied in some reports, increasing

the risk of type I statistical errors. Selective reporting, together with a large number of tested associations, small effect sizes, differences in design, definitions, outcomes and analytical approaches used, may produce spurious associations.⁵⁰ This may have biased the overall understanding of the role of environmental exposures on the development of asthma or wheeze in Latin-American children. A part of the observed selective reporting may be explained by publication bias, reflecting difficulties in publishing negative or non-conclusive findings and leading to selective reporting of positive results. However, recent provisions for online supplementary tables for most publishing platforms now allow authors to provide data and associations for all exposures measured.

Early-life infections have been shown to protect against atopy,¹⁵ but effects on asthma are still controversial. Evidence in this review points towards a higher risk of wheeze or asthma associated with early-life respiratory infections. Only five studies collected this information prospectively,^{9 25–27 45 46} and four of which reported an increased risk of wheeze or asthma following early-life respiratory infections.^{9 25 26 45 46} Respiratory syncytial virus bronchiolitis is considered to be an important risk factor for asthma,⁵¹ whereas rhinovirus has been associated with acute asthma exacerbations.^{52 53} These findings are difficult to interpret as most studies do not describe the type of respiratory infection or whether such infections were simply a manifestation of their underlying respiratory disease (transient wheeze or asthma). On the other hand, gastrointestinal and other chronic viral or bacterial diseases may not affect the risk of wheeze or asthma in Latin America.¹⁵

The association between intestinal parasites (mainly geohelminths) and asthma has been widely studied, and although a protective effect on atopy has been demonstrated,^{6 10 15} their effects on asthma remain unclear. An international meta-analysis showed no overall effect on asthma, though *Ascaris lumbricoides* was associated with an increased risk and hookworm with a decreased risk.⁵⁴ Similar findings can be seen in this review, with a predominantly protective effect of *Trichuris trichiura* on atopic wheeze and a higher risk of asthma or wheeze associated with *A. lumbricoides* infestation. The effect of intestinal helminths on asthma may depend on many factors, such as parasite species, intensity of infection, age of first infection and duration of infection.⁵⁵

Animals living around the home may increase the risk of infection with certain pathogens associated with asthma (eg, *Toxocara canis*).⁵⁵ Here, pet contact was not clearly associated with a higher risk of wheeze/asthma. A meta-analysis of international studies⁵⁶ found that dog exposures increased the risk of asthma slightly, whereas cat exposures reduced the risk. As furry animals may induce allergic diseases, it is difficult to ascertain whether they may increase the risk of asthma by increasing the risk of early-life infections or through their effect on atopy. Consistent protective effects across studies of contact with farm animals against asthma are one of the

most compelling observations in support of the hygiene hypothesis.⁵ This review provides only limited support for a protective role of such exposures in Latin-American populations.

Overcrowding, day-care attendance and having older siblings may increase the risk of early-life infections due to frequent and close contact with other children. However, there is no clear evidence of the effect these exposures have on childhood asthma.⁵⁷ In this review of Latin-American studies, these exposures in general were not associated with wheeze or asthma.

This review has several limitations. First, most of the studies included in the review were cross-sectional or case-control studies, which preclude establishing a time association between exposures and outcome. Second, the definition of asthma or wheeze differed between studies, complicating the analysis as not all wheeze is asthma, and although current wheeze is a good indicator of asthma for prevalence studies,² it may not be suitable for exploring asthma risk factors. More importantly the symptom 'wheeze' may be a manifestation of other respiratory pathologies, such as childhood respiratory infections that are a more frequent cause of chronic respiratory symptoms in Latin America than in other regions.¹² Within 'wheeze' may be included different disease processes with differing risk factors, as indicated by the observation from a recent meta-analysis of observational studies from industrialised countries that endotoxin exposure may increase the risk of wheeze in younger children but be protective against asthma in older children.⁵⁸ Asthma/wheeze likely encompasses a range of phenotypes and wide spectrum of disease severity associated with different patterns of risk factors. However, with the available data in this systematic review, it was not our aim to evaluate the effects of poor hygiene and infections on disease phenotypes or severity. Third, two-thirds of the studies were done in Brazil, with scarce representation of other large urban centres such as those present in Argentina, Peru or Uruguay. This may limit the generalisability of the findings to other Latin-American countries with different circumstances such as climate, socioeconomic level or diet. Finally, most studies did not provide results stratified by atopy, an important effect modifier. Previous studies have found contradictory effects of certain factors related to microbial exposure on either atopic or non-atopic asthma.⁵⁹

CONCLUSION

In conclusion, our findings in this systematic review do not settle the debate of whether the hygiene hypothesis is relevant or not to the high asthma prevalence in Latin-American children. Our analysis indicates a higher risk of wheeze and asthma in Latin America associated with acute respiratory infections in early life. Highly heterogeneous results regarding poor hygiene and early-life infections may be explained by difference in asthma phenotypes (atopic vs non-atopic) and control groups

used for comparison as well as different definitions used (current wheeze vs doctor's diagnosis of asthma). Selective reporting is common among observational studies exploring associations between environmental exposures and wheeze or asthma risk. Large prospective cohort studies with standardised outcomes are needed in Latin America to clarify the role of poor hygiene exposures and early-life infections on the development of childhood wheeze and asthma. Such studies should help guide policy makers on decisions of potential strategies to reduce the high asthma burden in Latin America.

Contributors CA-G, PJC and PG designed the study and protocol, interpreted the data and drafted the manuscript. CA-G extracted and summarised the data. All authors read and approved the final manuscript.

Funding This work was supported by the Wellcome Trust (grant 099938/B/12/Z) to CA-G and UK Aid for the benefit of developing countries (grant 5242) to PG.

Competing interests None declared.

Patient consent Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement All data generated or analysed during this study are included in this published article and its supplementary information files.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. Eder W, Ege MJ, von Mutius E. The asthma epidemic. *N Engl J Med* 2006;355:2226–35.
2. Masoli M, Fabian D, Holt S, et al. The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 2004;59:469–78.
3. Wills-Karp M, Santeliz J, Karp CL. The germless theory of allergic disease: revisiting the hygiene hypothesis. *Nat Rev Immunol* 2001;1:69–75.
4. Rook GA. Hygiene hypothesis and autoimmune diseases. *Clin Rev Allergy Immunol* 2012;42:5–15.
5. Riedler J, Braun-Fahrlander C, Eder W, et al. Exposure to farming in early life and development of asthma and allergy: a cross-sectional survey. *Lancet* 2001;358:1129–33.
6. Feary J, Britton J, Leonardi-Bee J. Atopy and current intestinal parasite infection: a systematic review and meta-analysis. *Allergy* 2011;66:569–78.
7. Schaub B, Lauener R, von Mutius E. The many faces of the hygiene hypothesis. *J Allergy Clin Immunol* 2006;117:969–77.
8. Weinmayr G, Weiland SK, Björkstén B, et al. Atopic sensitization and the international variation of asthma symptom prevalence in children. *Am J Respir Crit Care Med* 2007;176:565–74.
9. Barreto ML, Cunha SS, Fiaccone R, et al. Poverty, dirt, infections and non-atopic wheezing in children from a Brazilian urban center. *Respir Res* 2010;11:167.
10. Moncayo AL, Vaca M, Oviedo G, et al. Risk factors for atopic and non-atopic asthma in a rural area of Ecuador. *Thorax* 2010;65:409–16.
11. Pearce N, Ait-Khaled N, Beasley R, et al. Worldwide trends in the prevalence of asthma symptoms: phase III of the International Study of Asthma and Allergies in Childhood (ISAAC). *Thorax* 2007;62:758–66.
12. Mallol J, Solé D, Asher I, et al. Prevalence of asthma symptoms in Latin America: the International Study of Asthma and Allergies in Childhood (ISAAC). *Pediatr Pulmonol* 2000;30:439–44.
13. da Silva ER, Sly PD, de Pereira MU, et al. Intestinal helminth infestation is associated with increased bronchial responsiveness in children. *Pediatr Pulmonol* 2008;43:662–5.

14. Pereira MU, Sly PD, Pitrez PM, *et al.* Nonatopic asthma is associated with helminth infections and bronchiolitis in poor children. *Eur Respir J* 2007;29:1154–60.
15. Alcantara-Neves NM, Veiga RV, Dattoli VC, *et al.* The effect of single and multiple infections on atopy and wheezing in children. *J Allergy Clin Immunol* 2012;129:359–67.
16. Veiga RV, Cunha SS, Dattoli VC, *et al.* Chronic virus infections suppress atopy but not asthma in a set of children from a large Latin American city: a cross-section study. *BMC Pulm Med* 2011;11:24.
17. Asher MI, Keil U, Anderson HR, *et al.* International Study of Asthma and Allergies in Childhood (ISAAC): rationale and methods. *Eur Respir J* 1995;8:483–91.
18. ISAAC Phase One Manual. <http://isaac.auckland.ac.nz/phases/phaseone/phaseonemanual.pdf> (accessed Feb 2013).
19. Global Initiative for Asthma (GINA). *Global strategy for asthma management and prevention (2017 Update)*, 2017.
20. Alcantara-Neves NM, de S G Britto G, Veiga RV, *et al.* Effects of helminth co-infections on atopy, asthma and cytokine production in children living in a poor urban area in Latin America. *BMC Res Notes* 2014;7:817–0500-7-817.
21. Mendonça LR, Veiga RV, Dattoli VC, *et al.* Toxocara seropositivity, atopy and wheezing in children living in poor neighbourhoods in urban Latin America. *PLoS Negl Trop Dis* 2012;6:e1886.
22. Arrieta MC, Arévalo A, Stiemsma L, *et al.* Associations between infant fungal and bacterial dysbiosis and childhood atopic wheeze in a nonindustrialized setting. *J Allergy Clin Immunol* 2017.
23. Chatkin MN, Menezes AM, Victora CG, *et al.* High prevalence of asthma in preschool children in Southern Brazil: a population-based study. *Pediatr Pulmonol* 2003;35:296–301.
24. Chatkin MN, Menezes AM. [Prevalence and risk factors for asthma in schoolchildren in southern Brazil]. *J Pediatr* 2005;81:411–6.
25. Cooper PJ, Chico ME, Vaca MG, *et al.* Effect of Early-Life Geohelminth Infections on the Development of Wheezing at 5 Years of Age. *Am J Respir Crit Care Med* 2018;197:364–372.
26. Muñoz A, Menezes AM, Reichert FF, *et al.* [Wheezing phenotypes from birth to adolescence: a cohort study in Pelotas, Brazil, 1993–2004]. *J Bras Pneumol* 2008;34:347–55.
27. Zepeda G, Díaz P, Pinto R, *et al.* Seguimiento de lactantes hospitalizados por bronquiolitis por virus respiratorio sincicial: Evolución clínica, respuesta de atopia inflamatoria y marcadores. *Resultados preliminares. Revista chilena de enfermedades respiratorias* 2016;32:18–24.
28. Barreto BA, Sole D. Prevalence of asthma and associated factors in adolescents living in Belem (Amazon region), Para, Brazil. *Allergol Immunopathol* 2014;42:427–32.
29. Casagrande RR, Pastorino AC, Souza RG, *et al.* [Asthma prevalence and risk factors in schoolchildren of the city of São Paulo, Brazil]. *Rev Saude Publica* 2008;42:517–23.
30. Cooper PJ, Vaca M, Rodriguez A, *et al.* Hygiene, atopy and wheeze-eczema-rhinitis symptoms in schoolchildren from urban and rural Ecuador. *Thorax* 2014;69:232–9.
31. Moncayo AL, Vaca M, Oviedo G, *et al.* Effects of geohelminth infection and age on the associations between allergen-specific IgE, skin test reactivity and wheeze: a case-control study. *Clin Exp Allergy* 2013;43:60–72.
32. Azalim S, Camargos P, Alves AL, *et al.* Exposure to environmental factors and relationship to allergic rhinitis and/or asthma. *Ann Agric Environ Med* 2014;21:59–63.
33. Barraza Villarreal A, Sanín Aguirre LH, Téllez Rojo MM, *et al.* Risk factors for asthma in school children from Ciudad Juárez, Chihuahua. *J Asthma* 2003;40:413–23.
34. Rojas Molina N, Legorreta Soberanis J, Olvera Guerra F. [Prevalence and asthma risk factors in municipalities of the State of Guerrero, Mexico]. *Rev Alerg Mex* 2001;48:115–8.
35. Solis-Soto MT, Patiño A, Nowak D, *et al.* Association between environmental factors and current asthma, rhinoconjunctivitis and eczema symptoms in school-aged children from Oropeza Province-Bolivia: a cross-sectional study. *Environ Health* 2013;12:95–069X-12-95.
36. Beck AF, Huang B, Simmons JM, *et al.* Role of financial and social hardships in asthma racial disparities. *Pediatrics* 2014;133:431–9.
37. Dogaru CM, Nyffenegger D, Pescatore AM, *et al.* Breastfeeding and childhood asthma: systematic review and meta-analysis. *Am J Epidemiol* 2014;179:1153–67.
38. Kotaniemi-Syrjänen A, Vainionpää R, Reijonen TM, *et al.* Rhinovirus-induced wheezing in infancy-the first sign of childhood asthma? *J Allergy Clin Immunol* 2003;111:66–71.
39. Lim RH, Kobzik L, Dahl M. Risk for asthma in offspring of asthmatic mothers versus fathers: a meta-analysis. *PLoS One* 2010;5:e10134.
40. Mitchell EA, Beasley R, Keil U, *et al.* The association between tobacco and the risk of asthma, rhinoconjunctivitis and eczema in children and adolescents: analyses from Phase Three of the ISAAC programme. *Thorax* 2012;67:941–9.
41. Sigurs N, Gustafsson PM, Bjarnason R, *et al.* Severe respiratory syncytial virus bronchiolitis in infancy and asthma and allergy at age 13. *Am J Respir Crit Care Med* 2005;171:137–41.
42. Quiroz-Arcentales L, Hernández-Flórez LJ, Calderón CAA, *et al.* Enfermedad y síntomas respiratorios en niños de cinco municipios carboníferos del Cesar, Colombia. *Rev.salud pública* 2013;15:66–79.
43. Boneberger A, Haider D, Baer J, *et al.* Environmental risk factors in the first year of life and childhood asthma in the Central South of Chile. *J Asthma* 2011;48:464–9.
44. Del-Rio-Navarro B, Berber A, Blandón-Vijil V, *et al.* Identification of asthma risk factors in Mexico City in an International Study of Asthma and Allergy in Childhood survey. *Allergy Asthma Proc* 2006;27:325–33.
45. Brandão HV, Vieira GO, TdO V, *et al.* Fatores de risco socioeconômicos e ambientais associados à asma em crianças nascidas em maternidades públicas e privadas no Brasil. *Braz.j.allergy immunol* 2014;2:154–60.
46. Brandão HV, Vieira GO, Vieira TO, *et al.* Acute viral bronchiolitis and risk of asthma in schoolchildren: analysis of a Brazilian newborn cohort. *J Pediatr* 2017;93:223–9.
47. Coelho MA, de Pinho L, Marques PQ, *et al.* Prevalence and factors associated with asthma in students from Montes Claros, Minas Gerais, Brazil. *Cien Saude Colet* 2016;21:1207–16.
48. ISSAC Phase Two Study Modules. <http://isaac.auckland.ac.nz/phases/phasetwo/phasetwomodules.pdf> (accessed Feb 2013).
49. Phase III Environmental Questionnaire 13-14 years. http://isaac.auckland.ac.nz/phases/phasethree/environmentalquestionnaire/environmentalquestionnaire13_14.pdf (accessed Feb 2013).
50. Ioannidis JP. Why most published research findings are false. *PLoS Med* 2005;2:e124.
51. Szabo SM, Levy AR, Gooch KL, *et al.* Elevated risk of asthma after hospitalization for respiratory syncytial virus infection in infancy. *Paediatr Respir Rev* 2013;13(Suppl 2):S9–15.
52. Soto-Quiros M, Avila L, Platts-Mills TA, *et al.* High titers of IgE antibody to dust mite allergen and risk for wheezing among asthmatic children infected with rhinovirus. *J Allergy Clin Immunol* 2012;129:1499–505.
53. Heymann PW, Platts-Mills TA, Johnston SL. Role of viral infections, atopy and antiviral immunity in the etiology of wheezing exacerbations among children and young adults. *Pediatr Infect Dis J* 2005;24:S217–22.
54. Leonardi-Bee J, Pritchard D, Britton J. Asthma and current intestinal parasite infection: systematic review and meta-analysis. *Am J Respir Crit Care Med* 2006;174:514–23.
55. Cooper PJ. Toxocara canis infection: an important and neglected environmental risk factor for asthma? *Clin Exp Allergy* 2008;38:551–3.
56. Takkouche B, González-Barcala FJ, Etminan M, *et al.* Exposure to furry pets and the risk of asthma and allergic rhinitis: a meta-analysis. *Allergy* 2008;63:857–64.
57. Ramsey CD, Celedón JC. The hygiene hypothesis and asthma. *Curr Opin Pulm Med* 2005;11:14–20.
58. Mendy A, Gasana J, Vieira ER, *et al.* Endotoxin exposure and childhood wheeze and asthma: a meta-analysis of observational studies. *J Asthma* 2011;48:685–93.
59. Eduard W, Douwes J, Omenaas E, *et al.* Do farming exposures cause or prevent asthma? Results from a study of adult Norwegian farmers. *Thorax* 2004;59:381–6.
60. Kanobana K, Vereecken K, Junco Díaz R, *et al.* Toxocara seropositivity, atopy and asthma: a study in Cuban schoolchildren. *Trop Med Int Health* 2013;18:403–6.
61. van der Werff SD, Junco Díaz R, Reyneveld R, *et al.* Prediction of asthma by common risk factors: a follow-up study in Cuban schoolchildren. *J Investig Allergol Clin Immunol* 2013;23:415–20.
62. Wördemann M, Díaz RJ, Heredia LM, *et al.* Association of atopy, asthma, allergic rhinoconjunctivitis, atopic dermatitis and intestinal helminth infections in Cuban children. *Trop Med Int Health* 2008;13:180–6.
63. Bragagnoli G, Silva MT. Ascaris lumbricoides infection and parasite load are associated with asthma in children. *J Infect Dev Ctries* 2014;8:891–7.
64. Cooper PJ, Chico ME, Bland M, *et al.* Allergic symptoms, atopy, and geohelminth infections in a rural area of Ecuador. *Am J Respir Crit Care Med* 2003;168:313–7.
65. Endara P, Vaca M, Chico ME, *et al.* Long-term periodic anthelmintic treatments are associated with increased allergen skin reactivity. *Clin Exp Allergy* 2010;40:1669–77.

66. Freitas MS, Monteiro JC, Camelo-Nunes IC, *et al.* Prevalence of asthma symptoms and associated factors in schoolchildren from Brazilian Amazon islands. *J Asthma* 2012;49:600–5.
67. Garcia E, Aristizabal G, Vasquez C, *et al.* Prevalence of and factors associated with current asthma symptoms in school children aged 6–7 and 13–14 yr old in Bogotá, Colombia. *Pediatr Allergy Immunol* 2008;19:307–14.
68. Gomes de Luna MF, Gomes de Luna JR, Fisher GB, *et al.* Factors associated with asthma in adolescents in the city of Fortaleza, Brazil. *J Asthma* 2015;52:485–91.
69. Guimarães MAP, Fonseca MdeA, Amorim CR, *et al.* Sintomas asmáticos e fatores associados em crianças escolares. *Revista Baiana Saúde Pública* 2014;38:821–36.
70. Hagel I, Cabrera M, Hurtado MA, *et al.* Infection by *Ascaris lumbricoides* and bronchial hyper reactivity: an outstanding association in Venezuelan school children from endemic areas. *Acta Trop* 2007;103:231–41.
71. Han YY, Forno E, Badellino HA, *et al.* Antibiotic Use in Early Life, Rural Residence, and Allergic Diseases in Argentinean Children. *J Allergy Clin Immunol Pract* 2017;5:1112–8.
72. Han YY, Badellino HA, Forno E, *et al.* Rural residence, farming environment, and allergic diseases in Argentinean adolescents. *Pediatr Pulmonol* 2017;52:21–8.
73. Kuschner FC, Alves da Cunha AJ. Environmental and socio-demographic factors associated to asthma in adolescents in Rio de Janeiro, Brazil. *Pediatr Allergy Immunol* 2007;18:142–8.
74. Lima WL, Lima EV, Costa MR, *et al.* [Asthma and associated factors in students 13 and 14 years of age in São Luís, Maranhão State, Brazil]. *Cad Saude Publica* 2012;28:1046–56.
75. Maia JG, Marcopito LF, Amaral AN, *et al.* [Prevalence of asthma and asthma symptoms among 13 and 14-year-old schoolchildren, Brazil]. *Rev Saude Publica* 2004;38:292–9.
76. Palvo F, Toledo EC, Menin AM, *et al.* Risk factors of childhood asthma in Sao Jose do Rio Preto, Sao Paulo, Brazil. *J Trop Pediatr* 2008;54:253–7.
77. Prietsch SO, Fischer GB, César JA, *et al.* [Risk factors for recurrent wheezing in children under 13 years old in the South of Brazil]. *Rev Panam Salud Publica* 2006;20:331–7.
78. Ribeiro SA, Furuyama T, Schenkman S, *et al.* Atopy, passive smoking, respiratory infections and asthma among children from kindergarten and elementary school. *Sao Paulo Med J* 2002;120:109–12.
79. Silva MB, Amor AL, Santos LN, *et al.* Risk factors for *Toxocara* spp. seroprevalence and its association with atopy and asthma phenotypes in school-age children in a small town and semi-rural areas of Northeast Brazil. *Acta Trop* 2016.
80. Soto-Quiros M, Bustamante M, Gutierrez I, *et al.* The prevalence of childhood asthma in Costa Rica. *Clin Exp Allergy* 1994;24:1130–6.
81. Souza VM, Sales IR, Peixoto DM, *et al.* *Giardia lamblia* and respiratory allergies: a study of children from an urban area with a high incidence of protozoan infections. *J Pediatr* 2012;88:233–8.
82. Tintori SM, Carvalho deB, Dalva M, *et al.* Prevalência de asma e doenças alérgicas em escolares da faixa etária de 12 a 15 anos do município de Maringá-PR. *BBM rev. bras.med* 2013;70.
83. Toledo MF, Rozov T, Leone C. Prevalence of asthma and allergies in 13- to 14-year-old adolescents and the frequency of risk factors in carriers of current asthma in Taubaté, São Paulo, Brazil. *Allergol Immunopathol* 2011;39:284–90.
84. Cadore PS, Zhang L, Lemos LL, *et al.* Toxocariasis and childhood asthma: A case-control study. *J Asthma* 2016;53:601–6.
85. Jucá SC, Takano OA, Moraes LS, *et al.* [Asthma prevalence and risk factors in adolescents 13 to 14 years of age in Cuiabá, Mato Grosso State, Brazil]. *Cad Saude Publica* 2012;28:689–97.
86. López ML, Bojanich MV, Jacobacci JM, *et al.* [Toxocara canis and bronchial asthma]. *Medicina* 2010;70:75–8.
87. Mendoza DL, Socarrás SL, Sanabria MJB, *et al.* Asociación entre atopia, asma alérgica y anticuerpos IgE específicos para *Ascaris* en un grupo de niños de una ciudad de la Costa norte colombiana. *Salud Uninorte* 2008;24.
88. Moraes LS, Barros MD, Takano OA, *et al.* [Risk factors, clinical and laboratory aspects of asthma in children]. *J Pediatr* 2001;77:447–54.
89. Oliveira-Santos S, Motta-Franco J, Barreto I, *et al.* Asthma in adolescents – Prevalence trends and associated factors in northeast Brazil. *Allergol Immunopathol* 2015;43:429–35.
90. Rizzo MC, Naspitz CK, Fernández-Caldas E, *et al.* Endotoxin exposure and symptoms in asthmatic children. *Pediatr Allergy Immunol* 1997;8:121–6.