

# To intubate or not: ventilation is the question. A manikin-based observational study

Fatimata Seydou Sall,<sup>1,2</sup> Alban De Luca,<sup>1,2</sup> Lionel Pazart,<sup>1</sup> Aurore Pugin,<sup>1</sup> Gilles Capellier,<sup>2,3</sup> Abdo Khoury<sup>1,2</sup>

**To cite:** Sall FS, De Luca A, Pazart L, *et al.* To intubate or not: ventilation is the question. A manikin-based observational study. *BMJ Open Res* 2018;**5**:e000261. doi:10.1136/bmjresp-2017-000261

Received 9 November 2017  
Revised 20 June 2018

## ABSTRACT

**Introduction** There is a continuous debate concerning the superiority of endotracheal intubation on bag-valve-mask (BVM) ventilation in patients with cardiac arrest. In this manikin-based observational study, we evaluate and compare the performance of manual ventilation through a facemask (BVM) and an endotracheal tube (ETT).

**Methods** One hundred and forty healthcare providers were instructed to manually ventilate a manikin as they would do for a 75 kg adult patient in respiratory arrest. Each one was ventilating both through a facemask and an ETT for a 5 min period in a random order. Ventilatory parameters were measured by the ASL 5000 lung simulator and ventilation performance was analysed using a sliding window method published in a previous study to assess accurately ventilation efficiency.

**Results** The mean ventilation rate was high whatever the technique used (24 bpm). A weak relationship between manual ventilation performance and the type of interface used was observed ( $p=0.0484$ ). The overall rate of adequate ventilation was low even if we noticed a slight improvement when ventilating through an ETT (13.21% vs 7.5% of adequate ventilation). However, the rate of hyperventilation did not differ between mask and tube (79% vs 77%). A significant relationship is observed between professional category, the size of the hand squeezing the bag and manual ventilation performance ( $p<0.05$ ).

**Conclusion** Whatever the interface used, healthcare professionals are still struggling to perform manual ventilation efficiently according to international guidelines. Ventilation with an ETT does not prove to be significantly more efficient than with a facemask. It would be therefore important to recentre the debate on controlling ventilatory parameters with current devices. Focusing on training may maximise manual ventilation efficiency and minimise the loss of time during cardiopulmonary resuscitation.

## INTRODUCTION

Manual ventilation with facemask or endotracheal tube (ETT) interface (also called bag-valve ventilation) is frequently needed in many prehospital and in-hospital sceneries (during induction of general anaesthesia, after intubation while mechanical ventilation is being prepared, or when mechanical

## Key messages

- ▶ Whatever the interface used, healthcare professionals are still struggling to perform manual ventilation efficiently according to international guidelines.
- ▶ It would be therefore important to re-centre the debate on controlling ventilatory parameters with current devices and on training.

ventilation fails to achieve better patient oxygenation).<sup>1-3</sup> However, these basic and advanced airway management procedures (bag-valve-mask (BVM) with a facemask or endotracheal intubation (ETI) with an ETT) may have adverse events that can disturb patient haemodynamics and consequently impair survival.<sup>4,5</sup>

BVM is the most common method of ventilation used by emergency medical technicians. The challenge of this technique is to maintain patent airways and reduce leaks.<sup>6,7</sup> Several research studies have shown that healthcare professionals have trouble in ventilating adequately patients in accordance with international guidelines. This irregular and inadequate ventilation leads to either excessive airway pressure, gastric insufflation, and subsequent regurgitation and pulmonary aspiration or to hypercapnia and hypoxaemia.<sup>4,6-12</sup>

In opposition, ETI requires advanced skills in airway management and is only practised by experienced and highly trained healthcare professionals (mainly emergency physicians or anaesthetists).<sup>5,6,13</sup> However, this procedure might require time to perform and potentially lead to harmful events such as oesophageal and bronchial trauma, but also can sometimes be prolonged and delay the oxygenation of the patient or even completely fail.<sup>14,15</sup> Furthermore, it is important to minimise chest compression interruption during cardiopulmonary resuscitation in order to increase survival with



© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Inserm CIC 1431, University Hospital of Besançon, Besançon, France

<sup>2</sup>Department of Emergency Medicine and Critical Care, University Hospital of Besançon, Besançon, France

<sup>3</sup>Monash University, Melbourne, Victoria, Australia

## Correspondence to

Dr Fatimata Seydou Sall; fatimata.sarr@chu-besancon.fr

good neurological performances.<sup>16</sup> Thereby, while it is considered the 'gold standard technique' for advanced airway management, international guidelines recommend the use of laryngeal mask airway or BVM as alternatives for adults.<sup>13</sup>

There is a continuous debate regarding the superiority of ETI on BVM and their ability to provide adequate oxygenation to patients.<sup>17</sup> Indeed it remains important to know that, even if ETI is correctly performed, the main challenges of manual ventilation are to keep adequate and constant tidal volume and ventilation rate to prevent hyperinflation or hypoventilation. Therefore, we designed this prospective randomised bench study to compare the performance of manual ventilation with ETT and facemask, and assess the variability of the provided ventilation.

## MATERIALS AND METHODS

### Study design

In this study, we used the same protocol on BVM ventilation that we published in 2015<sup>18</sup> to compare the performance of BVM and ETI manual ventilation. This protocol was submitted for ethical approval but it was waived by the local ethics committee (CPP Est II). This manikin-based observational study was conducted in the Department of Emergency Medicine and Critical Care of Besançon. Briefly, 140 healthcare professionals, trained in basic life support during their initial training, were enrolled. They were still in service, aged over 18 years and coming from different independent structures (University Hospital of Besançon, Doubs Fire Department and Jussieu Ambulance Services). They were instructed to ventilate a manikin with two adult bag valves (Ambu and Laerdal bags) selected randomly in sealed envelopes, as they would do for a 75 kg adult patient in respiratory arrest. However, they were blinded regarding their ventilation performance transcribed on a computer. Ventilation was performed in protected airways with an ETT (8.0 mm ID). The size and the grip strength of the hand squeezing the bag were determined by using a tape measure and a Takei dynamometer, respectively. We define the level of professional experience based on their seniority at the moment of the trial enrolment. Thus, data of operator characteristics (professional category, grip strength of the hand keeping the bag and hand size) and ventilatory parameters (tidal volume ( $V_T$ ), ventilation frequency ( $V_R$ ), inspiratory and expiratory times (Itime and Etime)) were collected.

The manikin, a Laerdal Airway Management Trainer (Laerdal Medical, Stavanger, Norway), was connected to an ASL 5000 lung simulator (IngMar Medical, Pittsburgh, PA, USA) which simulated an apnoeic patient with a compliance of 70 mL/cmH<sub>2</sub>O and an airway resistance of 3.5 cmH<sub>2</sub>O/L/s. Ventilatory parameters (tidal volume, ventilation frequency, inspiratory and expiratory times) were measured by the lung simulator using the ASL SW V.3.3.106 software.

The manual ventilation performance was assessed using a new analysis method developed and published recently.<sup>18</sup> The algorithm of this method evaluates the performance of a 1 min window depending on tidal volume ( $V_T$ ) and ventilation rate ( $V_R$ ). It consists of segmenting every ventilation test into sliding windows of 1 min length each with a shift of three ventilation cycles each time. This will enable us to consider intraindividual, interindividual and time variability of ventilatory parameters. Regarding our simulated patient model (75 kg ideal body weight (IBW) with no respiratory pathology), we considered tidal volume from 300 to 600 mL and a ventilation rate between 8 and 15 bpm to be adequate. Thus, three situations were distinguished to assess the performance of a sliding window according to tidal volume ( $V_T$ ) and ventilation rate ( $V_R$ ) measurements during the 1 min period:

- Hyperventilation:  $V_R > 15$  bpm and/or mean  $V_T > 600$  mL.
- Adequate ventilation:  $V_R \leq 15$  bpm, mean  $V_T \leq 600$  mL with at least eight adequate ventilation cycles (with a  $V_T$  between 300 and 600 mL).
- Hypoventilation:  $V_R \leq 15$  bpm, mean  $V_T \leq 600$  mL and the number of adequate ventilation cycles  $< 8$ .

### Data analysis

Continuous data are expressed as mean  $\pm$  SD and as percentages for nominal variables. The comparison between professional categories and hand size and the different performance levels was done using a  $\chi^2$  test or Fisher exact test. OR, estimated by logistic regression, was used to analyse performance level for the multivariate model. A  $p < 0.05$  was considered statistically significant and SAS V.9.3 (SAS Institute) was used for statistical analysis and randomisation.

## RESULTS

Five hundred and sixty ventilation tests (280 for ventilation using appropriate facemask provided with each type of bag<sup>18</sup> and 280 for ventilation through an ETT) were performed by 140 healthcare professionals (45 physicians, 45 nurses and 50 rescuers) with a mean age of 37 (9) years. Among them, 45% (63/140) had a high level of professional experience and one-third were women. This study population is presented in table 1.

Results show that the mean ventilation rate was high (24 bpm) and similar when healthcare professionals ventilated through a facemask or an ETT. The mean tidal volume obtained with the ETT interface was greater than that obtained during manual ventilation with the facemask interface (462 vs 334 mL for ETT and facemask interfaces, respectively) (table 2).

The ventilation performance of healthcare professionals was also analysed and results show that manual ventilation performance was slightly influenced by the type of interface used ( $p = 0.0484$ ). The overall rate of adequate ventilation was low even if we noticed a minimal

**Table 1** Characteristics of study population (n=140)

Mean age $\pm$ SD (years)	37.28 $\pm$ 8.97
Sex, n (%)	
Female	47 (33.57)
Male	93 (66.43)
Professional category, n (%)	
Physicians	45 (32.14)
Emergency medicine physicians	29 (20.71)
Anaesthesia and critical care physicians	16 (11.43)
Nurses	45 (32.14)
Anaesthesia nurses	27 (19.29)
Emergency medicine nurses	18 (12.86)
First-aid workers	50 (35.71)
Firefighters	31 (22.14)
Emergency medical technicians	17 (12.14)
Red Cross first-aid rescuers	02 (1.43)
Professional experience, n (%)	
High ( $\geq$ 10 years)	63 (45.00)
Medium (5 $\leq$ n<10 years)	36 (25.70)
Little (<5 years)	41 (29.30)
Size of the hand squeezing the bag, n (%)	
Large ( $\geq$ 23 cm)	21 (15.00)
Medium (19 $\leq$ n<23 cm)	101 (72.14)
Small (15 $\leq$ n<19 cm)	18 (12.86)
Grip strength of the hand squeezing the bag, n (%)	
High ( $\geq$ 40 kgF)	53 (37.90)
Medium (20 $\leq$ n<40 kgF)	65 (46.40)
Weak (0 $\leq$ n<20 kgF)	22 (15.70)

improvement when ventilating through an ETT (13.21% vs 7.5% adequate ventilation tests). The rate of hyperventilation did not differ (79% vs 77%) (table 3).

We also evaluated some operator characteristics that may have an impact on ventilation performance. Multivariate analysis shows that manual ventilation performance was only impacted by the professional category ( $p=0.0002$  and  $p=0.0286$  for ETT and facemask

**Table 3** Ventilation performance analysis during the 5 min ventilation tests (n=280 tests for each ventilation technique)

Ventilation performance	Facemask (mean (%))	ETT (mean (%))
Adequate ventilation	21 (7.5)	37 (13.21)
Hypoventilation	37 (13.21)	27 (9.64)
Hyperventilation	222 (79.29)	216 (77.14)

ETT, endotracheal tube.

ventilations, respectively). The size of the hand squeezing the bag had a significant impact only on ETT ventilation performance ( $p=0.0175$  and  $p=0.5982$  for ETT and facemask, respectively). The other factors like grip strength, bag type (Ambu or Laerdal bag) and professional experience had no effect on ventilation performance ( $p>0.05$ ).

Figure 1 depicts more in detail the ventilation performance of the different professional categories and shows that physicians and nurses were more efficient than first-aid workers when ventilation was performed with an ETT (20%, 13% and 7%, respectively). This difference disappears when ventilation was performed with a facemask (9%, 6.67% and 7%, respectively, for physicians, nurses and first-aid workers). We observed also that all healthcare professionals hyperventilated whatever the interface used (>70%).


The impact of the hand size on ventilation performance is presented in figure 2. It shows that the larger the hand squeezing the bag was, the more excessive the ventilation was (90% vs 75% for large and medium hand size, respectively). When the hand was small, there were more hypoventilation tests compared with medium hand size results (16.67% vs 8.42%).

The intraindividual and interindividual variability of the two ventilation interface techniques is also analysed. There was not any intraindividual variability between facemask and ETT ventilations concerning ventilation rate ( $p>0.05$ ) while the variability of tidal volume was doubled with facemask compared with ETT ventilation ( $p<0.01$ ). However, a statistically significant difference was observed for the interindividual variability of tidal volumes between ETT and facemask interfaces (154 vs 132 mL,  $p=0.00001$ ) with a maximum mean tidal volume of 877 and 1004 mL and a minimum mean tidal volume

**Table 2** Ventilation parameter values measured during all the 5 min ventilation tests (n=280 tests for each ventilation technique)

Variable	Facemask			ETT		
	Mean $\pm$ SD	Lower quartile	Upper quartile	Mean $\pm$ SD	Lower quartile	Upper quartile
Ventilation rate ( $V_R$ , bpm)	24.09 $\pm$ 9.47	17.2	29.09	23.60 $\pm$ 9.66	16.27	29.09
Tidal volume ( $V_T$ , mL)	333.94 $\pm$ 124.19	245.6	419.95	462.31 $\pm$ 155.42	361.4	545.01
Inspiratory time (Itime, s)	0.76 $\pm$ 0.26	0.58	0.89	0.88 $\pm$ 0.34	0.66	1.03
Expiratory time (Etime, s)	2.10 $\pm$ 1.19	1.28	2.53	2.17 $\pm$ 1.31	1.27	2.63

ETT, endotracheal tube.




**Figure 1** Percentage of hyperventilation (black), adequate ventilation (grey) and hypoventilation (light grey) for professional categories (n=280 tests for each ventilation technique). ETT, endotracheal tube.

of 47 and 142 mL, respectively, for facemask and ETT ventilation tests.

## DISCUSSION

Ventilation is important for patient survival in cardiac arrest and/or respiratory distress. It is usually performed manually through a facemask or an ETT. These two ventilation techniques, mostly used by healthcare professionals in hospital and prehospital settings, have some problematic aspects.

ETI is considered to be the ‘gold standard’ because it maintains airway patency and avoids leaks, but we were interested to know whether, after the intubation, healthcare professionals ventilated adequately the patient in accordance with international guidelines. Indeed, inadequate ventilation is associated with impaired haemodynamics and increases morbidity and mortality. So we have evaluated and compared the performance of these



**Figure 2** Percentage of hyperventilation (black), adequate ventilation (grey) and hypoventilation (light grey) for hand size categories of the hand squeezing the bag (n=280 tests with an endotracheal tube (ETT)). Large hand size (≥23 cm), medium hand size (19 ≤ n < 23 cm) and small hand size (15 ≤ n < 19 cm).

two techniques and the factors involved, and the intra-individual and interindividual variability of the provided ventilation.

Healthcare professionals have performed 560 ventilation tests with an ETT (280) and a facemask (280). Among them, only 58 (10.4%) were adequate ventilation tests while the remaining 502 (89.6%) were inadequate ventilation tests (438 (78.2%) hyperventilation tests and 64 (11.4%) hypoventilation tests). This inadequate ventilation could increase the morbidity and the mortality of patients. Indeed, hyperventilation decreases the PaCO<sub>2</sub>, leading to central nervous system vasoconstriction, lower cerebral blood flow and secondary neurologic tissue injury, and can create free radicals and irreversible cellular damage.<sup>19–21</sup> Furthermore, the mean ventilation frequency was twice more than the adequate ventilation rate recommended by international guidelines.<sup>22–23</sup>

It was shown that hyperventilation is mostly caused by high ventilation rates rather than excessive volumes.<sup>24</sup> However, we observed a slight improvement in ventilation performance with the ETT than with the facemask (13% vs 7.5%), which was due to a reduction of hypoventilation sequences. This could be explained by the lack of leaks during ventilation with an ETT, leaks that remain difficult to control during ventilation with a facemask. Nonetheless, this rate of adequate ventilation stayed very low and proves that healthcare professionals are still struggling to perform manual ventilation efficiently and are hyperventilating in over 75% of cases whatever the interface used (79.3% vs 77.1%, respectively, for facemask and ETT ventilation tests).

In order to understand which factors may impact manual ventilation performance, we studied operator characteristics and showed that only the professional category and the size of the hand squeezing the bag affect manual ventilation performance. First-aid workers seem to have more difficulties to ventilate adequately in protected airway with 7% of adequate ventilation tests while 20% of physicians and 13% of nurses provided adequate ventilation. These results could be explained by the fact that first-aid workers, first on scene before advanced paramedics, are more accustomed to ventilate patients in unprotected airway. In these conditions, leaks could attenuate excessive ventilation volume provided to the patient. However, we observed that they also hyperventilated either with a facemask or with an ETT. This suggests that they tend to ventilate with an ETT as they use to do with a facemask (93% of inadequate ventilation tests for each technique). Only experienced and highly trained healthcare professionals such as advanced paramedics, physicians and nurses are authorised to secure the patient airways with an ETT and they are used to ventilate intubated patients, hence the high rate of efficient ventilation compared with first-aid workers (20% vs 7% and 13% vs 7% for physicians and nurses, respectively, vs first-aid workers). Conversely, physicians and nurses who performed efficient ventilation with ETT hypoventilated the lung with facemask interface. These results underline

that it would be necessary for healthcare professionals to adapt the appropriate approach to ventilate the patient when they use a facemask or an ETT interface and to take into account presence or absence of leaks.

Concerning the size of the hand squeezing the bag, it only has an influence when using an ETT. We observed that the participants with larger hand size (>23 cm) have major tendency to hyperventilate the patient, with 0% of adequate ventilation tests observed. When the hand size was medium (between 19 and 23 cm), there was a reduction of approximately 16% of hyperventilation. However, when the hand size is small, that is, under 19 cm, the percentage of hypoventilation increased by 8%. It might be helpful to raise awareness of healthcare professionals regarding the influence of these factors to adapt ventilation practices in order to optimise the ventilation efficiency (eg, healthcare professionals with large hands should squeeze the bag gently to avoid excessive ventilation). This may also explain why the interindividual variability between participants was so important. Indeed, we recorded mean tidal volumes varying from 47 to 877 mL with facemask and from 142 to 1004 mL with ETT. The same heterogeneous performances were observed on ventilation frequencies with mean  $V_R$  ranging from 8 to 53 bpm and from 8 to 48 bpm with facemask and ETT interfaces, respectively.

Although the debate continues in favour of tracheal intubation to secure the airway and maintain patency despite a lack of clear evidence,<sup>16 25</sup> intubating the patient may lead to several complications both at the time, during and after intubation, and even result in loss of effectiveness of resuscitation.<sup>26–28</sup> A study of Hanif *et al* has shown that this technique was associated with decreased survival to hospital discharge of adult non-traumatic patients compared with ventilation with a facemask.<sup>29</sup> So, if ventilation with an ETT does not prove to be significantly more efficient than with a facemask, as we showed in this study, it would be important to refocus the debate on controlling ventilatory parameters and developing high-performance medical tools that can maximise ventilation efficiency and minimise the loss of time during cardiopulmonary resuscitation. Indeed, immediate and early cardiopulmonary resuscitation with a short interruption of chest compressions is vital to increase patient survival. These medical tools can be real-time feedback devices that would instantly indicate to the healthcare professionals what they actually do during manual ventilation.

This study presents some limits as it is a manikin study and we still cannot confirm that the ventilation performance described herein is a reliable representation of what could be obtained in clinical trials. In real clinical situations, especially in intubated patients, measuring  $\text{ETCO}_2$  by capnography can help the healthcare professional to better ventilate. However, most of studies dealing with ventilation performance in the literature are manikin-based studies.<sup>30</sup> Leaks in manikin ventilation (BVM) are more common than in real use on patients.

Only few masks work effectively with a manikin, and are usually not supplied with the bags. But leaks induce users to ventilate faster hoping to ensure adequate ventilation. In addition, we decided to evaluate ventilation performance and not the difficulty related to intubation technique. This latter issue has been well documented in the literature and therefore was not considered in this study. Thus, the manikin was intubated before the 280 ventilation tests were performed. Furthermore, we still cannot confirm that the BVM ventilation performance described in this study and obtained with a new analysis method is a reliable representation of what could be obtained in clinical trials and in other centres.

In summary, the high failure rate of manual ventilation observed in this study shows that ‘to intubate or not to intubate the patient’ is not really the question, especially when its benefits are still being debated. ETI is certainly considered the ‘gold standard technique’ but unfortunately, ventilation is still poorly performed afterwards inducing other complications in addition to those associated with intubation. New devices as the Fastrach or laryngeal mask may be good alternative to increase the rate of adequate ventilation. We think we should focus more on how to improve manual ventilation efficiency with current devices and minimise these complications. We should also consider how to teach healthcare professionals to adapt their ventilation practices according to the airway management technique used.

**Acknowledgements** We thank all the staff of the Departments of Emergency Medicine and of Anaesthesia and Critical Care of Besançon University Hospital, the staff of the Doubs Fire Department and Besançon Jussieu Ambulances teams who voluntarily participated in this research project and made it possible.

**Contributors** FSS carried out the bibliography, drafted the protocol and the manuscript. ADL defined the new analysis method and realised bench tests. AP performed the statistical analysis and drafted the statistical section. LP participated in the conception and design of the study. GC guided the work and corrected the manuscript. AK supervised the work and corrected the manuscript. All authors approved the final manuscript.

**Funding** This work is supported by an unrestricted grant from the European Commission (FEDER) in Franche-Comté, Bpi France, the Greater Besançon Urban Area Community (CAGB), the Regional Council of Franche-Comté and the General Council of Doubs Department.

**Competing interests** None declared.

**Patient consent** Obtained.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** No additional data are available.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## REFERENCES

1. Turki M, Young MP, Wagers SS, *et al*. Peak pressures during manual ventilation. *Respir Care* 2005;50:340–4.
2. Clayton TJ, Pittman JA, Gabbott DA. A comparison of two techniques for manual ventilation of the lungs by non-anaesthetists: the bag-valve-facemask and the cuffed oropharyngeal airway (COPA) apparatus. *Anaesthesia* 2001;56:756–9.

3. Patman S, Jenkins S, Stiller K. Manual hyperinflation--effects on respiratory parameters. *Physiother Res Int* 2000;5:157-71.
4. Gammon RB, Shin MS, Groves RH, et al. Clinical risk factors for pulmonary barotrauma: a multivariate analysis. *Am J Respir Crit Care Med* 1995;152(4 Pt 1):1235-40.
5. Alexander R, Hodgson P, Lomax D, et al. A comparison of the laryngeal mask airway and Guedel airway, bag and facemask for manual ventilation following formal training. *Anaesthesia* 1993;48:231-4.
6. Doerges V, Sauer C, Ocker H, et al. Airway management during cardiopulmonary resuscitation--a comparative study of bag-valve-mask, laryngeal mask airway and combitube in a bench model. *Resuscitation* 1999;41:63-9.
7. Nolan JP, Soar J, Zideman DA, et al. Adult basic life support and use of automated external defibrillators: European Resuscitation Council Guidelines for Resuscitation 2010. *Resuscitation* 2010;81:1219-76.
8. Stone BJ, Chantler PJ, Baskett PJ. The incidence of regurgitation during cardiopulmonary resuscitation: a comparison between the bag valve mask and laryngeal mask airway. *Resuscitation* 1998;38:3-6.
9. Wenzel V, Keller C, Idris AH, et al. Effects of smaller tidal volumes during basic life support ventilation in patients with respiratory arrest: good ventilation, less risk? *Resuscitation* 1999;43:25-9.
10. von Goedecke A, Voelckel WG, Wenzel V, et al. Mechanical versus manual ventilation via a face mask during the induction of anesthesia: a prospective, randomized, crossover study. *Anesth Analg* 2004;98:260-3.
11. Handley AJ, Koster R, Monsieurs K, et al. European Resuscitation Council guidelines for resuscitation 2005. Section 2. Adult basic life support and use of automated external defibrillators. *Resuscitation* 2005;67(Suppl 1):S7-23.
12. Lim JS, Cho YC, Kwon OY, et al. Precise minute ventilation delivery using a bag-valve mask and audible feedback. *Am J Emerg Med* 2012;30:1068-71.
13. Boyle M, Flavell E. Which is more effective for ventilation in the prehospital setting during cardiopulmonary resuscitation, the laryngeal mask airway or the bag-valve-mask? A review of the literature. *J Emerg Prim Health Care* 2012;8:2.
14. Benoit JL, Prince DK, Wang HE. Mechanisms linking advanced airway management and cardiac arrest outcomes. *Resuscitation* 2015;93:124-7.
15. Divatia JV, Bhowmick K. Complications of endotracheal intubation and other airway management procedures. *Indian J Anaesth* 2005;49:308-18.
16. Neumar RW, Otto CW, Link MS, et al. Part 8: adult advanced cardiovascular life support: 2010 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2010;122(18 Suppl 3):S729-67.
17. Henlin T, Michalek P, Tyll T, et al. Oxygenation, ventilation, and airway management in out-of-hospital cardiac arrest: a review. *Biomed Res Int* 2014;2014:1-11.
18. Khoury A, Sall FS, De Luca A, et al. Evaluation of bag-valve-mask ventilation in manikin studies: what are the current limitations? *Biomed Res Int* 2016;2016:1-8.
19. Lurie KG, Yannopoulos D, McKnite SH, et al. Comparison of a 10-breaths-per-minute versus a 2-breaths-per-minute strategy during cardiopulmonary resuscitation in a porcine model of cardiac arrest. *Respir Care* 2008;53:862-70.
20. Aufderheide TP, Sigurdsson G, Pirralo RG, et al. Hyperventilation-induced hypotension during cardiopulmonary resuscitation. *Circulation* 2004;109:1960-5.
21. Gaither JB, Spaite DW, Bobrow BJ, et al. Balancing the potential risks and benefits of out-of-hospital intubation in traumatic brain injury: the intubation/hyperventilation effect. *Ann Emerg Med* 2012;60:732-6.
22. Perkins GD, Travers AH, Berg RA, et al. Part 3: adult basic life support and automated external defibrillation: 2015 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2015;95:e43-69.
23. Kleinman ME, Brennan EE, Goldberger ZD, et al. Part 5: Adult Basic Life Support and Cardiopulmonary Resuscitation Quality: 2015 American Heart Association guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2015;132(18 Suppl 2):S414-35.
24. O'Neill JF, Deakin CD. Do we hyperventilate cardiac arrest patients? *Resuscitation* 2007;73:82-5.
25. Tiah L, Kajino K, Alsakaf O, et al. Does pre-hospital endotracheal intubation improve survival in adults with non-traumatic out-of-hospital cardiac arrest? A systematic review. *West J Emerg Med* 2014;15:749-57.
26. Wang HE, Yealy DM. Out-of-hospital endotracheal intubation: where are we? *Ann Emerg Med* 2006;47:532-41.
27. Scott DB. Endotracheal intubation: friend or foe. *BMJ* 1986;292:157-8.
28. Timmermann A, Russo SG, Eich C, et al. The out-of-hospital esophageal and endobronchial intubations performed by emergency physicians. *Anesth Analg* 2007;104:619-23.
29. Hanif MA, Kaji AH, Niemann JT. Advanced airway management does not improve outcome of out-of-hospital cardiac arrest. *Acad Emerg Med* 2010;17:926-31.
30. Ward PA, Irwin MG. Man vs. manikin revisited - the ethical boundaries of simulating difficult airways in patients. *Anaesthesia* 2016;71:1399-403.