

Electronic Table 1 Logistic regression analysis for risk factors associated with in-hospital deaths in \*AE-IPF.

	<sup>†</sup> OR	95% <sup>‡</sup> CI	<i>P</i> value
Age (year)	0.96	0.87–1.07	0.46
<sup>‡</sup> PaO <sub>2</sub> / <sup>§</sup> FiO <sub>2</sub>	1.00	0.99–1.01	0.87
**WBC (×10 <sup>3</sup> /μL)	1.38	1.04–1.83	0.03
<sup>††</sup> Hb (mg/dL)	0.51	0.30–0.85	0.01
<sup>‡‡</sup> CRP (mg/dL)	1.00	0.90–1.13	0.94
Antibiotic therapy	4.89	0.39–60.56	0.22

\*AE-IPF: acute exacerbation of idiopathic pulmonary fibrosis. <sup>†</sup>OR: odds ratio. <sup>‡</sup>CI: confidence interval. <sup>‡</sup>PaO<sub>2</sub>: partial pressure of arterial oxygen. <sup>§</sup>FiO<sub>2</sub>: fraction of inspired oxygen. \*\*WBC: white blood cell count. <sup>††</sup>Hb: haemoglobin. <sup>‡‡</sup>CRP: C-reactive protein.