The proportion of males increased as severity increased. The percentage of males who had no, mild, moderate or severe OSA was 64.6%, 72.0%, 86.0 and 87.0%, respectively.

Mean age was significantly lower ($P<0.001$) in the no OSA compared to all 3 OSA categories. Those with moderate OSA were also significantly older ($P<0.05$) than those with mild OSA.

Those with severe OSA had a significantly higher average BMI compared to all other categories ($P<0.05$).

Average neck circumference significantly increased with severity of OSA.

There was no significant difference in average ESS between groups.

Conclusion Seventy-five percent of pre-operative patients with a STOP-BANG score ≥ 3 had some degree of OSA. The severity of OSA was dependent on gender, age, BMI and neck circumference. ESS was not helpful in predicting presence or severity of OSA.

P021 NON-INVASIVE VENTILATION FOR OBESE PATIENTS WITH OSA FAILING REGULAR CPAP: A PROSPECTIVE OBSERVATIONAL COHORT STUDY

^{1,2}Athanasios Ishak*, ²Michelle Ramsay, ^{1,2}Nicholas Hart, ^{1,2}Joerg Steier. ¹King's College London Faculty of Life Sciences and Medicine, Centre for Human and Applied Physiological Science (CHAPS), London, UK; ²Guy's and St Thomas' NHS Foundation Trust Lane Fox Respiratory Unit, London, UK

10.1136/bmjresp-2019-bssconf.21

Introduction Continuous positive airway pressure (CPAP) therapy is the standard treatment for obstructive sleep apnoea (OSA), however, patients frequently have sub-optimal long-term adherence. We hypothesised that bilevel positive airway pressure (BPAP) therapy may improve adherence and outcomes in patients who cannot tolerate CPAP.

Methods Patients with OSA who had sub-optimal CPAP usage (<4 hours/day) and were referred to our sleep centre between 2014–2017 for BPAP were included. Age, gender, body-mass-index (BMI), co-morbidities, CPAP use and reasons for failure, Epworth Sleepiness Score (ESS), sleep study data, spirometry data and average maximum nightly compliance were recorded.

Results We included 52 patients with OSA who required CPAP >15 cmH₂O (71% male, age 58 (15) years, BMI 42.6 (10.1) kg/m², AHI 51.1 (30.4)/hour); 62% had respiratory co-morbidities affecting nocturnal breathing (COPD, OHS). CPAP was used for 199 (106–477) days prior to referral. Reasons for CPAP failure were intolerance of pressures (23%), uncontrolled symptoms (23%), mask problems (21%), adverse effects (13%), claustrophobia (8%), co-morbidities (8%), and other issues (4%). Lower expiratory positive airway pressures (EPAP) were needed to control nocturnal breathing compared to CPAP (10 (8–12) vs 16.8 (15.7–19.2) cmH₂O, $p=0.001$) and patients achieved better adherence (7.0 (4.0–8.5) vs 2.5 (1.6–6.7) hours/night, $p=0.028$) and better symptom control (ESS 4.0 (1.0–7.0) vs 10.0 (6.0–17.0) points, $p=0.039$) on BPAP.

Discussion In patients with OSA with limited success on CPAP therapy, BPAP is better tolerated and achieves sufficient respiratory and symptom control.

P022 AN INVESTIGATION INTO THE EFFECTS OF A 20-MINUTE NAP-OPPORTUNITY ON COGNITIVE FUNCTION IN STUDENTS AGED 18–30

¹Suraiya Haddad, ¹Rezwana Rahman, ¹Felix Wong*, ¹W Jackson, ^{1,2}Mary Morrell. ¹Imperial College London, London, UK; ²Academic Unit of Sleep and Ventilation, National Heart and Lung Institute, Imperial College London, Royal Brompton and Harefield National Health Service Foundation Trust

10.1136/bmjresp-2019-bssconf.22

Introduction Sleep-deprivation is increasing in prevalence and compensatory napping is becoming common practice. Little is known about the cognitive benefits derived from a nap in sleep-deprived individuals, or within the student population. Furthermore, few studies have investigated the cognitive benefits of napping using the incongruent Stroop test. This study aimed to investigate the effects of a nap-opportunity on cognitive function in individuals aged 18–30.

Methods A randomised, crossover study was conducted. Participants were recruited from Imperial College London and were allocated to either control (no nap-opportunity) or intervention (20-minute nap-opportunity) groups.

The incongruent Stroop test was used to measure reaction time (RT) and percentage accuracy, which were used as markers of cognitive function. Δ RT and Δ percentage accuracy, between control and nap-opportunity groups were compared. Paired and unpaired t-tests were used for the analysis of RT. Wilcoxon signed-rank and Mann-Whitney-U tests were used to test the percentage accuracy. The study was approved by the Imperial College Medical Education Ethics Committee.

Results Participants ($n=29$) had a mean age of 20 ($SD\pm 1.4$) years, 55% male. In those who were acutely sleep-deprived, the RT improved in the intervention group following a 20-minute nap-opportunity, compared to the control group ($p=0.01$; figure 1). Overall, no significant difference in Δ

Abstract P022 Figure 1 Dot plot showing reaction times in incongruent Stroop test, measured before and after a 20 minute nap-opportunity in individuals who slept for <8 hrs the night before the test ($n=18$) and $=8$ hrs the night before the test ($n=11$). Data presented as mean $\pm 95\%$ CI. Paired t-test was used to compare reaction times before and after nap-opportunity. Unpaired t-tests were used to compare baselines and endpoints of the two groups. Significant change in reaction time was observed following nap-opportunity in individuals who slept <8 Hours before the test day ($p=0.01275$) * $=P<0.05$, ** $=P<0.01$, *** $=P<0.001$