

Discussion This study suggests that a ‘one-size fits all’ approach to mattresses may not be appropriate and contrasting body types need different levels of support to improve overall spinal alignment. The use of simple anthropometric measurements could make the selection of the most appropriate mattress easier for the public.

P003

RANDOMISED, PLACEBO-CONTROLLED STUDY OF SOLRIAMFETOL FOR EXCESSIVE DAYTIME SLEEPINESS IN NARCOLEPSY TYPES 1/2

^{1,2}Jed Black*, ³Yves Dauvilliers, ⁴Giuseppe Plazzi, ⁵Colin M Shapiro, ^{6,7}Geert Mayer, ⁸Bruce C Corser, ⁹Helene A Emsellem, ¹Dan Chen, ^{1,10}Lawrence P Carter, ¹¹Hao Wang, ¹²Michael J Thorpy. ¹Jazz Pharmaceuticals, Palo Alto, USA; ²Stanford Center for Sleep Sciences and Medicine, Palo Alto, USA; ³Gui-de-Chauliac Hospital, CHU Montpellier, Montpellier, France; ⁴University of Bologna, Bologna, Italy; ⁵University of Toronto, Toronto, Canada; ⁶Hephata Klinik, Schwalmstadt, Germany; ⁷Philipps University Marburg, Marburg, Germany; ⁸Sleep Management Institute, Cincinnati, USA; ⁹The Center for Sleep and Wake Disorders, Chevy Chase, USA; ¹⁰University of Arkansas for Medical Sciences, Little Rock, USA; ¹¹Kite Pharma, Inc., Santa Monica, USA, (previously from Jazz Pharmaceuticals, Palo Alto, CA, USA); ¹²Albert Einstein College of Medicine, Bronx, USA

10.1136/bmjresp-2019-bssconf.3

Introduction Solriamfetol (formerly JZP-110), a dopamine and norepinephrine reuptake inhibitor, has been approved in the United States to improve wakefulness in adults with excessive daytime sleepiness (EDS) associated with narcolepsy (75–150 mg) or obstructive sleep apnoea (OSA; 37.5–150 mg). A Marketing Authorisation Application for these indications is under review with the European Medicines Agency. This phase 3 study assessed safety and efficacy of solriamfetol in participants with narcolepsy types 1 and 2 (NT1/2).¹

Methods In this 12-week, double-blind, randomised, placebo-controlled study, participants with or without cataplexy were randomised to solriamfetol 75 mg, 150 mg, 300 mg, or placebo. Eligibility criteria: NT1/2 diagnosis; mean sleep latency <25 minutes on Maintenance of Wakefulness Test (MWT); Epworth Sleepiness Scale (ESS) score ≥10. Exclusion criteria: medications that could affect EDS or cataplexy; night-time or variable shift work; other conditions causing EDS.

Results 236 participants received ≥1 dose of solriamfetol (67.2% female; 80.2% white). Baseline MWT mean sleep latency: 7.5 minutes; baseline ESS score: 17.2. Solriamfetol significantly increased MWT sleep latency at week 12 ($P<0.0001$ for 300 mg and 150 mg); least squares (LS) mean change: 12.3 minutes for 300 mg, 9.8 for 150 mg, 4.7 for 75 mg, and 2.1 for placebo. Solriamfetol significantly decreased ESS scores at week 12 ($P<0.0001$ 150 mg and 300 mg; $P<0.05$ 75 mg). LS mean change in ESS: -6.4 for 300 mg, -5.4 for 150 mg, -3.8 for 75 mg, and -1.6 for placebo. Most common treatment-emergent adverse events (TEAEs; ≥5%): headache, nausea, decreased appetite, nasopharyngitis, dry mouth, and anxiety. Discontinuations due to TEAEs were more frequent in solriamfetol 150 mg and 300 mg groups.

Discussion Solriamfetol improved wakefulness and reduced EDS in participants with NT1/2. Most AEs were mild to moderate.

Support Jazz Pharmaceuticals.

REFERENCE

1. Thorpy MJ, et al. *Ann Neurol* 2019;**85**(3):359–370.

P004

SOLRIAMFETOL FOR EXCESSIVE DAYTIME SLEEPINESS IN OBSTRUCTIVE SLEEP APNOEA: A RANDOMISED CONTROLLED TRIAL

¹Paula K Schweitzer*, ^{2,3}Geert Mayer, ^{4,5}Russell Rosenberg, ^{6,7}Gary K Zammit, ⁸Mark Gotfried, ⁹Dan Chen, ^{9,10}Lawrence P Carter, ¹¹Hao Wang, ^{8,12}Jed Black, ¹³Atul Malhotra, ¹⁴Kingman P Strohl. ¹Sleep Medicine and Research Center, St. Luke's Hospital, Chesterfield, USA; ²Hephata Klinik, Schwalmstadt, Germany; ³Philipps University Marburg, Marburg, Germany; ⁴NeuroTrialsResearch, Inc., Atlanta, USA; ⁵Atlanta School of Sleep Medicine, Atlanta, USA; ⁶Clinilabs Drug Development Corporation, New York, USA; ⁷Icahn School of Medicine at Mount Sinai, New York, USA; ⁸Pulmonary Associates, Phoenix, USA; ⁹Jazz Pharmaceuticals, Palo Alto, USA; ¹⁰University of Arkansas for Medical Sciences, Little Rock, USA; ¹¹Kite Pharma, Inc., Santa Monica, USA, (previously from Jazz Pharmaceuticals, Palo Alto, CA, USA); ¹²Stanford Center for Sleep Sciences and Medicine, Palo Alto, USA; ¹³Division of Pulmonary, Critical Care and Sleep Medicine, University of California, San Diego, USA; ¹⁴Case Western Reserve University, Cleveland, USA

10.1136/bmjresp-2019-bssconf.4

Introduction Obstructive sleep apnoea (OSA) is often associated with persistent excessive daytime sleepiness (EDS) despite sleep apnoea therapy. There are currently no approved treatments in the European Union for the treatment of EDS in this population. Solriamfetol (formerly JZP-110), a dopamine and norepinephrine reuptake inhibitor, has been approved in the United States to improve wakefulness in adults with EDS associated with narcolepsy (75–150 mg) or OSA (37.5–150 mg). A Marketing Authorisation Application for these indications is under review with the European Medicines Agency. This study evaluated the efficacy and safety of solriamfetol for treatment of EDS in participants with OSA with current or prior sleep apnoea treatment.¹

Methods In this double-blind, placebo-controlled, parallel-group phase 3 trial, participants with OSA and associated EDS were randomly assigned to solriamfetol 37.5 mg, 75 mg, 150 mg, or 300 mg or placebo for 12 weeks.

Results Of 476 randomised participants, 459 were included in the prespecified efficacy analyses. Co-primary endpoints (Maintenance of Wakefulness Test sleep latency, Epworth Sleepiness Scale score) were met at all solriamfetol doses ($P<0.05$), with dose-dependent effects observed at week 1 and maintained over the study duration. All doses except 37.5 mg resulted in significantly higher percentages of participants reporting improvement on Patient Global Impression of Change (key secondary endpoint; $P<0.05$). Adverse events (AEs) were reported in 47.9% of placebo- and 67.9% of solriamfetol-treated participants; 5 participants experienced serious AEs (2 [1.7%] placebo, 3 [0.8%] solriamfetol); none were deemed related to study drug. The most common AEs with solriamfetol were headache (10.1%), nausea (7.9%), decreased appetite (7.6%), anxiety (7.0%), and nasopharyngitis (5.1%).

Discussion Solriamfetol significantly improved wakefulness and reduced sleepiness in participants with OSA and EDS. Most AEs were mild or moderate.

Support Jazz Pharmaceuticals.

REFERENCE

1. Schweitzer PK, et al. *Am J Respir Crit Care Med* 2019;**199**(11):1421–1431.