

**Supplemental Table 3: Associations between radiographic indices of disease severity at the time of diagnosis and symptoms**

Radiographic Indices of Disease Severity <sup>a</sup>	No. Assessed	Symptoms						
		Respiratory			Constitutional			Both <sup>b</sup>
		Dry Cough n (%)	Prod. Cough n (%)	Hemoptysis n (%)	Fever n (%)	Night Sweats n (%)	Weight Loss n (%)	
<b>Cavitation</b>								
Yes	103	65 (63.1)	24 (23.3)	30 (29.1)	45 (43.6)	25 (24.2)	57 (55.3)	74 (71.8)
No	210	84 (40.0)	26 (12.4)	16 (7.6)	56 (26.6)	27 (12.9)	64 (30.4)	77 (36.6)
<b>Bilateral</b>								
Yes	118	69 (58.4)	22 (18.6)	27 (22.8)	47 (39.8)	22 (18.6)	59 (50.0)	70 (59.3)
No	195	80 (41.0)	28 (14.4)	19 (9.7)	54 (27.6)	30 (15.4)	62 (31.7)	81 (41.5)
<b>Moderate/ Far-Advanced</b>								
Yes	159	97 (61.0)	36 (22.6)	37 (23.2)	67 (42.1)	37 (23.2)	82 (51.5)	106 (66.6)
No	154	52 (33.7)	14 (9.1)	9 (5.8)	34 (22.1)	15 (9.8)	39 (25.3)	45 (29.2)

<sup>a</sup>Radiographic indices are defined in the text. Those without ‘bilateral’ disease could have either unilateral disease, or no disease.

<sup>b</sup>‘Both’ refers to any combination of respiratory and constitutional symptoms