

Online supplement

**Efficacy of recombinant thrombomodulin for poor prognostic cases of  
acute exacerbation in idiopathic interstitial pneumonia: Secondary  
analysis of the SETUP trial**

Table S1

**Table S1.** Adverse events and cause of death according to the prognostic staging in rhTM arm

	Poor prognostic stage (n=23)	Good prognostic stage (n=16)	p-value <sup>§</sup>
<b>Adverse events</b>	5	2	0.678
Bleeding	2*	0	0.503
Others			
Infection	1	0	
Hepatitis	1	0	
DVT	1	0	
Unspecified vascular event <sup>#</sup>	0	1	
Cerebral infarction	0	1	
<b>Dead cases</b>	11	2	0.037
Causes of death**			
AE	10	1	
Infection	1	0	
Unspecified vascular event <sup>#</sup>	0	1	

Abbreviations: rhTM, recombinant human soluble thrombomodulin; DVT, deep vein thrombosis; AE, acute exacerbation.

§: Frequency of adverse events and dead cases between poor and good prognostic stage in rhTM arm was compared by Fisher's exact test.

\*: Nasal bleeding and bleeding from gastric ulcer

\*\* : None of the dead cases underwent autopsy.

#: Back pain occurred 38 days after the commencement of thTM and the patient died on the same day. Dissection of aorta was suspected; however, a postmortem examination was not performed.