





# Prognostic significance of serum cytokines during acute exacerbation of idiopathic interstitial pneumonias treated with thrombomodulin

Toru Arai <sup>1</sup>, Hiroto Matsuoka,<sup>2</sup> Masaki Hirose,<sup>1</sup> Hiroshi Kida,<sup>3,4</sup> Suguru Yamamoto,<sup>5</sup> Yoshitaka Ogata,<sup>6</sup> Masahide Mori,<sup>3</sup> Kazuyoshi Hatsuda,<sup>1</sup> Chikatoshi Sugimoto,<sup>1</sup> Kazunobu Tachibana,<sup>1</sup> Masanori Akira,<sup>1</sup> Yoshikazu Inoue <sup>1</sup>

**To cite:** Arai T, Matsuoka H, Hirose M, *et al*. Prognostic significance of serum cytokines during acute exacerbation of idiopathic interstitial pneumonias treated with thrombomodulin. *BMJ Open Resp Res* 2021;**8**:e000889. doi:10.1136/bmjresp-2021-000889

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjresp-2021-000889>).

Received 1 February 2021  
Accepted 11 July 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

**Correspondence to**  
Dr Yoshikazu Inoue;  
[giichiyi@me.com](mailto:giichiyi@me.com)

## ABSTRACT

**Background** Acute exacerbation (AE) has been reported to herald a poor prognosis in idiopathic pulmonary fibrosis and is now thought to do so in idiopathic interstitial pneumonias (IIPs). However, the pathophysiology of AE-IIPs is not sufficiently understood. In our previously reported SETUP trial, we found better survival in patients with AE-IIPs treated with corticosteroids and thrombomodulin than in those treated with corticosteroids alone. In that study, we collected serum samples to evaluate changes in cytokine levels and retrospectively examined the prognostic significance and pathophysiological role of serum cytokines in patients with AE-IIPs.

**Methods** This study included 28 patients from the SETUP trial for whom serial serum samples had been prospectively obtained. AE-IIPs were diagnosed using the Japanese Respiratory Society criteria. All patients were treated with intravenous thrombomodulin and corticosteroids from 2014 to 2016. Serum levels of 27 cytokines were measured using Bio-Plex. The high-resolution CT pattern at the time of diagnosis of AE was classified as diffuse or non-diffuse.

**Results** Univariate analysis revealed that higher serum levels of interleukin (IL)-2, IL-7, IL-9, IL-12, IL13, basic fibroblast growth factor, granulocyte-macrophage colony-stimulating factor, interferon- $\gamma$  inducible protein-10, platelet-derived growth factor and regulated on activation, normal T cell expressed and secreted (RANTES) at AE were significant predictors of 90-day survival. The HRCT pattern was also a significant clinical predictor of 90-day survival. Multivariate analysis with stepwise selection identified a higher serum RANTES level at AE to be a significant predictor of 90-day survival, including after adjustment for HRCT pattern. Multivariate analysis with stepwise selection suggested that a marked increase in the serum IL-10 level on day 8 could predict 90-day mortality.

**Conclusions** A higher serum RANTES level at AE the time of diagnosis predicted a good survival outcome, and an elevated serum IL-10 level on day 8 predicted a poor survival outcome.

**Trial registration number** UMIN000014969.

## Key messages

- What is the most important cytokine to predict survival of acute exacerbation (AE) in idiopathic interstitial pneumonias (IIPs)?
- We found that a higher serum regulated on activation, normal T cell expressed and secreted level at the time of diagnosis of AE predicted a good survival outcome, and an elevated serum interleukin-10 level on day 8 predicted a poor outcome.
- The result might suggest pathophysiological significance of these cytokines and lead to the new treatment of AE-IIPs.

## INTRODUCTION

Idiopathic pulmonary fibrosis (IPF) is a lung disease of unknown aetiology and has a poor prognosis.<sup>1,2</sup> Some patients with IPF experience rapid potentially fatal deterioration<sup>1-4</sup> known as acute exacerbation (AE) of IPF (AE-IPF). AE is the most common cause of death in IPF,<sup>4</sup> despite affected patients usually being treated with corticosteroids according to the 2011 American Thoracic Society (ATS)/European Respiratory Society (ERS)/Japanese Respiratory Society (JRS)/Latin American Thoracic Association guidelines for diagnosis and management of IPF.<sup>1</sup> Therefore, a better understanding of the pathophysiology of AE-IPF is required to improve its management. Furthermore, the prognosis of AE in unclassifiable idiopathic interstitial pneumonias (IIPs) or interstitial lung disease other than IIPs is as poor as that of IPF.<sup>5,6</sup>

Recombinant human soluble thrombomodulin (rhTM), which contains only the extracellular domain of thrombomodulin, has recently been approved for treatment of disseminated intravascular coagulation (DIC)<sup>7</sup> and has been reported to be effective



for AE-IPF<sup>8</sup> when coupled with conventional corticosteroid therapy. We have previously reported on the efficacy of rhTM in patients with AE-IIPs who participated in our prospective multicentre SETUP trial.<sup>9</sup> Thrombomodulin regulates coagulation<sup>10</sup> and inhibits inflammation by preventing adhesion of neutrophils to the endothelium, decreasing the expression of adhesion molecules and enhancing the barrier function of endothelium and degradation of high-mobility group box 1.

Although the pathophysiology of AE-IPF is unknown, disordered integrity of epithelial cells, cellular inflammation, cytokines, matrix metalloproteinases, coagulation and fibrinolysis are likely to be involved, and rapid changes in these phenomena are thought to contribute to AE-IPF.<sup>4</sup> Collard *et al*<sup>11</sup> found that serum Krebs von den Lungen (KL)-6, interleukin (IL)-6 and thrombomodulin levels were higher in patients with AE-IPF than in those with stable IPF. Moreover, using a small number of lung specimens from patients with AE-IPF or stable IPF, Konishi *et al*<sup>12</sup> found that gene expression of AE-IPF was associated with more severe epithelial injury and proliferation.

Although there have been a few reports on changes in serum levels of several cytokines in patients with AE-IPF,<sup>13–15</sup> their importance in the pathophysiology and prognosis of AE-IPF and AE-IIPs has not been examined in sufficient depth. Therefore, in this retrospective study, we evaluated the prognostic and pathophysiological significance of serum cytokine levels in patients with AE-IIPs who had participated in the SETUP trial and from whom serum samples had been collected.

## MATERIALS AND METHODS

### Subjects

The SETUP trial included 39 patients with a diagnosis of AE-IIPs based on the modified JRS criteria<sup>16</sup> who had been prospectively enrolled and treated with rhTM and conventional therapy, including corticosteroids and immunosuppressants, from 2014 to 2016. Patients with severe non-pulmonary disease, life-threatening bleeding, clinically significant infection, known hypersensitivity to rhTM, a cerebrovascular event within the past year or surgery within the previous month were excluded. Pregnancy or the possibility of pregnancy was an additional exclusion criterion. Serum samples had been collected from 28 of the 39 patients at the time of diagnosis of AE-IIPs (on day 1) and from 22 patients on day 8 (online supplemental figure 1). Patient characteristics of 28 patients with serum samples in the present study were similar to the rest of the subjects in SETUP trial (n=11) without serum samples (online supplemental table 1).

The trial was registered with the University Hospital Medical Information Network Center (28 August 2014). All participants in SETUP provided written informed consent.

## METHODS

### Diagnosis of underlying IIPs and AE-IIPs

The underlying IPF was diagnosed based on the ATS/ERS/JRS/Latin American Thoracic Association guidelines for diagnosis and management of IPF.<sup>1</sup> The diagnosis of IIPs was based on the ATS/ERS/JRS statement,<sup>17</sup> and IIPs that could not be diagnosed as IPF was classified as non-IPF IIPs. Patients with non-IPF IIPs did not undergo surgical lung biopsy and were diagnosed to have unclassifiable IIPs.

AE-IIPs were diagnosed based on the following modified JRS diagnostic criteria for AE-IPF.<sup>16</sup> (1) Within 1 month, the following three conditions should be satisfied: (A) progressively worsening dyspnoea; (B) new ground-glass opacities evident on high-resolution CT (HRCT) scans superimposed on background reticular opacity, traction bronchiectasis, traction bronchiolectasis or honeycombing; and (C) a reduction in resting partial pressure of oxygen in arterial blood (PaO<sub>2</sub>) of >10 Torr compared with previous measurements. (2) Other causes of acute respiratory impairment, such as infection, pneumothorax, cancer, pulmonary embolism and congestive cardiac failure, should be excluded. However, the possibility that some patients had AE triggered by infection<sup>4</sup> could not be excluded because bronchoalveolar lavage could not be performed in all cases.

### Treatment for AE-IIPs

Details of the treatment of the patients with AE-IIPs in the SETUP trial are provided in online supplemental table 2. Treatment included prednisolone with/without an immunosuppressant after intravenous administration of high-dose methylprednisolone for three consecutive days. Polymyxin B-immobilised fibre column direct hemoperfusion (PMX/DHP) therapy<sup>13 15</sup> was administered in one case using the Toraymyxin system (Toray Medical, Tokyo, Japan), which was attached for 4–6 hours at a flow rate of 80 mL/min and repeated once within 48 hours. Nafamostat mesilate (Torii Pharmaceuticals, Tokyo, Japan) was used to reduce the risk of clot formation in the catheter. Anticoagulant and antiplatelet drugs were permitted only to treat comorbidities. Invasive or non-invasive positive pressure ventilation or nasal high flow therapy was introduced to maintain oxygenation in some patients with severe respiratory failure. rhTM was started at a rate of 380 U/kg/day for 6 days, as approved for treatment of DIC,<sup>7–9</sup> on the day of AE-IIPs diagnosis.

### Evaluation of AE-IIPs

AE-IIPs were evaluated before and at the time of diagnosis of AE as described in our previous report.<sup>9</sup> Briefly, the severity of IIPs (stages I–IV, mildest to most severe) in the stable state was evaluated in most cases within 6 months before the time of diagnosis of AE according to the JRS criteria.<sup>16</sup> Shortness of breath was evaluated based on the modified Medical Research Council score.<sup>18</sup> DIC was diagnosed according to the Japanese Association for Acute Medicine

scoring system.<sup>19</sup> The fibrin degradation product level was classified as higher ( $\geq 10$  mg/L) or lower ( $< 10$  mg/L). The HRCT pattern at the time of diagnosis of AE-IIPs was classified as peripheral, multifocal or diffuse using the definition proposed by Akira *et al*<sup>3</sup> and subsequently according to whether it was diffuse or non-diffuse. Patients with a PaO<sub>2</sub>/fraction of inspired oxygen (FiO<sub>2</sub>) ratio  $\leq 200$ <sup>20</sup> were deemed to have severely impaired oxygenation.

### Evaluation of serum biomarkers

Serum KL-6 levels were measured using a commercial ELISA kit (Eizai, Tokyo, Japan). The KL-6 cut-off level was 500 U/mL.<sup>9</sup> Cytokine levels in the serum samples (pg/mL) were quantified using the Bio-Plex Suspension Array System with Bio-Plex Pro Human Cytokine Group 27-Plex Panel (Bio-Rad Laboratories, Hercules, California, USA) according to the manufacturer's instructions.<sup>15</sup> This panel was selected because we could simultaneously measure serum levels of chemokines, fibrotic cytokines and angiogenic cytokines, and it was used in our previous study.<sup>15</sup> The cytokines measured included IL-1 $\beta$ , IL-1ra, IL-2, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12 (p70), IL-13, IL-15, IL-17A, eotaxin/CC chemokine ligand (CCL) 11, basic fibroblast growth factor (b-FGF), granulocyte colony-stimulating factor (G-CSF), granulocyte-macrophage colony-stimulating factor, interferon (IFN)- $\gamma$ , IFN- $\gamma$  inducible protein 10 (IP-10)/CXC chemokine ligand (CXCL) 10, monocyte chemotactic protein-1 (MCP-1)/CCL2, macrophage inflammatory protein (MIP)-1 $\alpha$ /CCL3, MIP-1 $\beta$ /CCL4, platelet-derived growth factor (PDGF)-BB, regulated on activation, normal T cell expressed and secreted (RANTES)/CCL5, tumour necrosis factor (TNF)- $\alpha$  and vascular endothelial growth factor (VEGF)-A. The cytokine levels in each patient, out of the range below or above, were defined to the upper range or half of the lower range, respectively. Serum IL-1 $\beta$ , IL-15 and VEGF-A levels at the time of diagnosis of AE were out of range in more than half of all cases (n=26, n=22 and n=19, respectively). These three cytokines were excluded from further examination.

### Statistical analysis

Continuous and categorical data are presented as the median (IQR) or the number (percentage) and were compared between the rhTM and control arms using the Wilcoxon rank-sum test and Fisher's exact test, respectively. Logistic regression analysis was performed to predict 90-day survival using clinical parameters, serum cytokine levels at the time of diagnosis of AE and the increase in serum cytokine levels on day 8. All statistical analyses were performed using JMP software V.11 (SAS Institute). All p values are two sided. A p value  $< 0.05$  was considered statistically significant.

## RESULTS

### Patient characteristics

The 28 patients with AE-IIPs comprised eight with IPF and 20 with non-IPF IIPs. Clinical parameters before

and at the time of diagnosis of AE were similar between those with AE-IPF and those with AE-non-IPF. The 90-day survival rate was similar between these two groups (table 1). Underlying IIPs were treated with antifibrotic drugs (n=5; nintedanib for one case and pirfenidone for four cases) and prednisolone (n=9) before the time of diagnosis of AE (table 1). Univariate logistic regression analysis revealed that use of antifibrotic drugs in a stable state, a diffuse HRCT pattern and higher serum KL-6 and lactate dehydrogenase levels at the time of diagnosis of AE were significant clinical predictors of 90-day mortality (table 2). Use of antifibrotic drugs in a stable state and a diffuse HRCT pattern at the time of diagnosis of AE were significant predictors in multivariate analysis. Use of antifibrotic drugs at stable state was significantly associated with introduction of long-term oxygen therapy (p=0.0047) and disease severity stage IV (p=0.0047) before the diagnosis of AE (online supplemental table 3).

There was no significant difference in serum cytokine levels between the AE-IPF and AE-non-IPF groups (online supplemental table 4).

### Survival and cytokines at the time of diagnosis of AE

Serum cytokine levels were compared between patients who were alive at 90 days and those who were not. Serum IL-2, IL-7, IL-9, IL-13, GM-CSF, IP-10, PDGF-BB and RANTES/CCL5 levels were higher in survivors than in non-survivors (table 3). Univariate logistic regression analysis revealed higher IL-2, IL-7, IL-9, IL-12, IL13, b-FGF, GM-CSF, IP-10, PDGF-BB and RANTES/CCL5 levels to be significant predictors of 90-day survival (table 4). Multivariate logistic analysis with stepwise selection revealed that IL-7 and RANTES/CCL5 should be included in the model; multivariate analysis using these two parameters identified RANTES/CCL5 to be a significant predictor of 90-survival (table 4). RANTES/CCL5 was a significant predictor after adjustment for HRCT pattern at the time of diagnosis of AE and antifibrotic drug use before AE (table 4). RANTES/CCL5 was a significant predictor of 90-day survival in AE-IIPs without use of antifibrotic drugs before AE (p=0.010, data not shown).

### RANTES/CCL5 level and clinical parameters

The serum RANTES/CCL5 level was not significantly associated with any clinical parameter suggesting severe underlying IIPs in a stable state or during AE, including use of long-term oxygen therapy in a stable state and the PaO<sub>2</sub>/FiO<sub>2</sub> and HRCT pattern at the time of diagnosis of AE. There was a significant correlation between the serum RANTES/CCL5 level and administration of antifibrotic drugs before AE (online supplemental table 5); however, only five patients were treated with these agents (table 1).



**Table 1** Patient characteristics in this study\*

Parameters	AE-IIPs (n=28)	AE-IPF (n=8)	AE-non-IPF (n=20)	P value
<b>Stable state</b>				
Sex, male/female	21/7	8/0	13/7	0.0749
Smoking, yes/no	19/9	7/1	12/8	0.2144
mMRC, ≤1/≥2	10/18	3/5	7/13	1.0000
Stage, I–III/IV	17/11	4/4	13/7	0.6715
LTOT before AE, yes/no	11/17	4/4	7/13	0.6175
Prednisolone before AE, yes/no	9/19	2/6	7/13	1.0000
Antifibrotic drugs, yes/no	5†/23	2‡/6	3§/17	0.6056
<b>At the time of diagnosis of AE</b>				
Age, years	74.5 (69.75–79.0)	74.5 (69.5–78.25)	74.5 (69.75–84.0)	0.7405
HRCT pattern, diffuse/non-diffuse	12/16	4/4	8/12	0.6908
PaO <sub>2</sub> /FiO <sub>2</sub> ratio, ≤200/>200	17/11	6/2	11/9	0.4188
WCC, /μL	10 150 (9050–12 950)	10 300 (9000–14 400)	10 150 (9250–12 950)	0.9594
LDH, U/L	317.0 (245.5–404.5)	296.5 (246.5–414.25)	341.0 (245.5–403.75)	0.9797
KL-6, U/mL	1196 (889–2,142)	1024.5 (889.5–1394)	1346 (849.75–2359)	0.3091
CRP, mg/dL	12.38 (5.07–14.28)	5.455 (0.38–12.56)	2.982 (0.871–9.815)	0.8787
FDP, 10</>≥10 mg/L	21/7	4/4	17/3	0.1423
90-day survival, yes/no	19/9	6/2	13/7	1.0000

\*Each parameter was compared between the AE-IPF and AE-non-IPF groups using Fisher's exact test or the Wilcoxon rank-sum test.

†Pirfenidone (n=4) and nintedanib (n=1).

‡Pirfenidone (n=1) and nintedanib (n=1).

§Pirfenidone (n=3) and nintedanib (n=0).

AE, acute exacerbation; CRP, C reactive protein; FDP, fibrin degradation product; FiO<sub>2</sub>, fraction of inspired oxygen; HRCT, high-resolution CT; IIPs, idiopathic interstitial pneumonias; IPF, idiopathic pulmonary fibrosis; KL-6, Krebs von den Lungen-6; LDH, lactate dehydrogenase; LTOT, long-term oxygen therapy; mMRC, modified medical research council score; PaO<sub>2</sub>, arterial oxygen tension; WCC, white blood cell count.

### Cytokine levels from days 1–8

By day 8, there was a significant increase in serum IL-4, IL-13, eotaxin and b-FGF levels and a significant decrease in serum IL1-ra, IL-9, IL-10, IFN-γ, IP-10, MIP-1β, RANTES/CCL5 and TNF-α levels (online supplemental table 6).

### Survival and increased cytokines on day 8

The increase in serum IL-2, IL-5, IL-7, IL-9, IL-10, G-CSF, IFN-γ and MCP-1 levels on day 8 was significantly smaller in survivors than in non-survivors at 90 days (table 5). The median increase in the levels of these cytokines except for IL-2 was less than zero, suggesting a decrease on day 8 in 90-day survivors. Univariate logistic regression analysis revealed that an increase in IL-7, IL-10, G-CSF, IFN-γ and MCP-1 levels suggested mortality at 90 days, and multivariate analysis with stepwise selection suggested that an increase in IL-10 on day 8 could predict 90-day mortality (table 6).

### Increased serum IL-10 and clinical parameters

There was no association between an increased IL-10 level on day 8 and any other clinical parameter at the time of diagnosis of AE or in a stable state (online supplemental table 7).

### DISCUSSION

We have examined the levels of multiple cytokines in serum samples that were prospectively collected from patients with AE-IIPs enrolled in the SETUP trial.<sup>9</sup> This type of evaluation is rare. We found that a higher serum RANTES/CCL5 level at the time of diagnosis of AE-IIPs suggested a good prognosis after treatment with corticosteroids and thrombomodulin. Furthermore, a greater increase in serum IL-10 at 1 week after the time of diagnosis of AE was a significant predictor of a poor prognosis.

RANTES/CCL5 is reportedly a chemotactic factor for lymphocytes, eosinophils, monocytes and mast cells and is produced by lymphocytes, fibroblasts, platelets, endothelial cells and epithelial cells.<sup>21</sup> RANTES/CCL5 has been found to be elevated in the bronchoalveolar lavage fluid of patients with nonspecific interstitial pneumonia<sup>22</sup> and in those with IPF.<sup>23</sup> Pierce *et al*<sup>24</sup> showed that RANTES/CCL5 works with CCL21 to affect pulmonary fibrosis. RANTES/CCL5 is reported to increase inflammation and fibrosis in the presence of renal or hepatic disease.<sup>25</sup> Hence, the good prognosis suggested by a higher RANTES/CCL5 level at the time of diagnosis of AE in our study indicates that it has an important pathophysiological role in AE-IIPs.

**Table 2** Prognostic significance of clinical parameters in patients with AE-IIPs determined by logistic regression analysis

Parameter	OR*	95% CI	P value
<b>Univariate</b>			
<b>Stable state</b>			
Sex, male/female	1.2499	0.2060 to 10.2093	0.8140
Smoking, yes/no	2.2400	0.4135 to 12.3435	0.7376
Diagnosis, IPF/non-IPF	0.6190	0.0765 to 3.5887	0.6037
mMRC, ≤1/≥2	1.1666	0.2251 to 6.9237	0.8560
Stage, I-III/IV	2.7083	0.5354 to 14.8724	0.2275
LTOT before AE, yes/no	2.7083	0.5354 to 14.8724	0.2275
Prednisolone before AE, yes/no	2.2400	0.4135 to 12.3435	0.3431
Antifibrotic drugs before AE, yes/no	14.400	1.6803 to 319.0268	0.0137
<b>At the time of diagnosis of AE</b>			
Age, years	0.9929	0.8995 to 1.0986	0.8863
HRCT pattern, diffuse/non-diffuse	9.8000	1.7196 to 83.0325	0.0091
PaO <sub>2</sub> /FiO <sub>2</sub> ratio, ≤200/>200	0.7291	0.1425 to 3.8110	0.7014
WCC, /μL	1.0001	0.9999 to 1.0004	0.1607
LDH, U/L	1.0065	1.0007 to 1.0145	0.0244
KL-6, U/mL	1.0009	1.0001 to 1.0020	0.0146
CRP, mg/dL	0.9678	0.8582 to 1.0777	0.6268
FDP, ≥10/<10 mg/L	4.2666	0.7166 to 28.9076	0.1102
<b>Multivariate†</b>			
HRCT pattern, diffuse/non-diffuse	13.3634	1.7053 to 285.2057	0.0118
Antifibrotic drugs before AE, yes/no	21.3180	1.6243 to 860.5019	0.0179

\*An OR >1 means each categorical parameter or increase in each continuous parameter indicates a high risk of 90-day mortality.

†Multivariate analysis with a stepwise selection procedure was performed to predict 90-day survival using significant parameters with a p value <0.05.

AE, acute exacerbation; CRP, C reactive protein; FDP, fibrin degradation product; FiO<sub>2</sub>, fraction of inspired oxygen; HRCT, high-resolution CT; IIPs, idiopathic interstitial pneumonias; IPF, idiopathic pulmonary fibrosis; KL-6, Krebs von den Lungen-6; LDH, lactate dehydrogenase; LTOT, long-term oxygen therapy; mMRC, modified Medical Research Council score; PaO<sub>2</sub>, arterial oxygen tension; WCC, white blood cell count.

We also found that serum IL-7, IL-9, IL-13, GM-CSF, IP-10, PDGF-BB and RANTES/CCL5 levels were significantly higher in survivors than in non-survivors. PDGF<sup>26</sup> and IL-13<sup>27</sup> are profibrotic cytokines, similar to RANTES/CCL5,<sup>21–25</sup> while IL-7,<sup>28,29</sup> GM-CSF<sup>30</sup> and IP-10/CXCL10<sup>31</sup> are antifibrotic cytokines. Whether IL-9 is profibrotic<sup>32</sup> or antifibrotic<sup>33</sup> is controversial; however, IL-9 is associated with regulation of pulmonary fibrosis. The results of our study suggest that an early increase in fibrosis-associated profibrotic and antifibrotic cytokines might affect the prognosis of AE-IIPs.

The pathophysiology of AE-IIPs is not yet fully clarified; however, it is thought to involve acute epithelial injury superimposed on chronic progressive interstitial lung disease. If the epithelial cells regenerate rapidly, the intact alveolar epithelial cell layer can suppress proliferation of fibroblasts and pulmonary fibrosis may be inhibited.<sup>26,34,35</sup> Although overgrowth of fibroblasts and overproduction of collagen are not favourable for pulmonary fibrosis, keratinocyte growth factor secreted by fibroblasts helps recovery from injury-stimulated proliferation of alveolar epithelial cells.<sup>36</sup> Hence, rapid

recruitment of fibroblasts at the site of injury by fibrosis-related cytokines and increased production of keratinocyte growth factor might lead to rapid repair of alveolar epithelial cells. Rapid reduction in fibrosis-related cytokine levels and proliferating fibroblasts might be necessary to prevent development of fibrosis.

The pathophysiology of AE-IPF has some similarity to that of the COVID-19 caused by SARS-CoV-2.<sup>37</sup> The virus infects the epithelium of the upper and lower airways via ACE 2 receptors, and diffuse alveolar damage occurs after extensive apoptosis of alveolar epithelial cells. Recently, Sugiyama *et al*<sup>37</sup> reported a low CCL17 level at the time of diagnosis of SARS-CoV-2 infection to be a significant predictor of severe COVID-19. CCL17 is reported to be a profibrotic cytokine in mice with bleomycin-induced lung injury<sup>38</sup> and to be elevated in bronchioalveolar lavage fluid from patients with IPF.<sup>39</sup> The disadvantage of lower levels of profibrotic cytokines in early lung injury might be similar between COVID-19 infection and AE-IPF or AE-IIPs.

Fibrocytes are circulating cells with fibroblast-like properties that are involved in both normal tissue wound

**Table 3** Serum cytokines at the time of diagnosis of AE-IIPs: comparison between 90-day survivors and non-survivors\*

Parameters	Survivors (n=19)	Non-survivors (n=9)	P value
IL-ra	95.48 (25.88 to 248.77)	161.6 (6.76 to 766.6)	0.8240
IL-2	0.430 (0.340 to 0.950)	0.240 (0.155 to 0.405)	0.0128
IL-4	0.300 (0.190 to 0.440)	0.270 (0.205 to 0.335)	0.4163
IL-5	1.380 (1.010 to 2.320)	1.010 (0.780 to 1.615)	0.0847
IL-6	0.480 (0.24 to 1.390)	1.100 (0.190 to 2.150)	0.7490
IL-7	1.410 (1.080 to 1.970)	0.800 (0.445 to 1.340)	0.0138
IL-8	3.510 (2.410 to 7.140)	3.970 (2.110 to 27.220)	0.9021
IL-9	2.150 (1.940 to 3.160)	1.790 (1.600 to 2.045)	0.0058
IL-10	0.900 (0.590 to 1.610)	0.590 (0.375 to 1.775)	0.3370
IL-12	0.260 (0.120 to 0.490)	0.120 (0.050 to 0.175)	0.0738
IL-13	0.210 (0.120 to 0.380)	0.070 (0.040 to 0.105)	0.0083
IL-17	1.090 (0.087 to 1.720)	1.090 (0.700 to 1.335)	0.5059
Eotaxin/CCL11	17.55 (10.17 to 32.92)	18.79 (10.98 to 20.76)	0.6055
b-FGF	5.860 (4.720 to 7.850)	4.720 (4.410 to 6.025)	0.0823
G-CSF	5.210 (0.980 to 25.140)	1.560 (0.34 to 22.688)	0.4151
GM-CSF	0.070 (0.040 to 0.320)	0.020 (0.020 to 0.040)	0.0119
IFN $\gamma$	0.800 (0.530 to 2.220)	0.620 (0.510 to 1.410)	0.5218
IP-10/CXCL10	605.6 (307.6 to 1293.8)	215.4 (154.8 to 538.7)	0.0195
MCP-1/CCL2	5.650 (3.280 to 22.530)	4.500 (2.640 to 13.145)	0.4029
MIP-1 $\alpha$ /CCL3	0.690 (0.400–1.230)	0.510 (0.280 to 8.205)	0.3758
PDGF-BB	87.81 (68.72 to 144.2)	47.88 (21.35 to 91.49)	0.0366
MIP-1 $\beta$ /CCL4	248.2 (186.1 to 307.1)	200.3 (193.6 to 355.3)	0.6759
RANTES/CCL5	2370 (1803 to 5610)	1568 (1333 to 1859)	0.0085
TNF- $\alpha$	4.680 (4.040 to 6.420)	3.840 (3.205 to 5.130)	0.0518

\*Serum cytokine levels at the time of diagnosis of AE (pg/mL) were compared between 90-day survivors and non-survivors using the Wilcoxon rank-sum test.

G-CSF; granulocyte colony-stimulating factor; AE, acute exacerbation; b-FGF, basic fibroblast growth factor; CCL, CC chemokine ligand; CXCL, CXC chemokine ligand; GM-CSF, granulocyte-macrophage colony-stimulating factor; IFN, interferon; IIPs, idiopathic interstitial pneumonias; IL, interleukin; IP-10, IFN- $\gamma$  inducible protein; MCP, monocyte chemotactic protein; MIP, macrophage inflammatory protein; PDGF, platelet-derived growth factor; RANTES, regulated on activation, normal T cell expressed and secreted; TNF, tumour necrosis factor.

repair and fibrosis. Fibrocytes secrete a variety of cytokines to augment fibrotic lesions by activating resident fibroblasts.<sup>40</sup> Fibrocytes are an important source of fibroblasts and myofibroblasts. In one study, circulating fibrocytes indicated a poor prognosis in IPF,<sup>41</sup> with circulating fibrocyte counts in the blood of patients with AE-IPF found to be elevated by up to 10-fold in comparison with the values in stable IPF.

Human fibrocytes express several chemokine receptors, including CCR2, CCR5, CCR7 and CXCR4.<sup>42</sup> RANTES/CCL5 is known to induce migration of circulating fibrocytes in patients with severe asthma<sup>43</sup> and might be associated with pulmonary fibrosis. Although this hypothesis supports the notion that RANTES/CCL5 worsens pulmonary fibrosis, Tai *et al*<sup>44</sup> demonstrated in a mouse model that intravenous infusion of fibrocytes ameliorated lipopolysaccharide-induced acute lung injury by decreasing inflammatory cytokine and chemokine levels and reducing accumulation of neutrophils in

the lung. Early recruitment of an appropriate number of fibrocytes might lead to normal repair instead of pulmonary fibrosis; this hypothesis is consistent with our finding that higher levels of serum RANTES/CCL5 at the time of diagnosis of AE are associated with a good prognosis in patients with AE-IIPs.

IL-7 inhibits production of transforming growth factor (TGF)- $\beta$  and fibroblast signalling.<sup>28</sup> Moreover, IL-7 and TGF- $\beta$  play counter-regulatory roles in synthesis of collagen by fibroblasts in pulmonary fibrosis.<sup>29</sup> We have shown that a higher IL-7 level at the time of diagnosis of AE can predict a good prognosis and that the IL-7 level was significantly lower on day 8 than at the time of diagnosis. We have previously reported that an increase in IL-7 24–48 hours after the start of PMX/DHP therapy predicts a good survival outcome.<sup>15</sup> These results may seem inconsistent; however, the exact time course of the serum IL-7 level from the time of diagnosis of AE-IIPs (day 1) to day 8 is not known, and the peak level may be on day 3.

**Table 4** Prognostic significance of serum cytokines for AE-IIPs determined by logistic regression analysis for 90-day mortality

Parameters	OR*	95% CI	P value
Univariate analysis			
IL-ra	1.0002	0.9986 to 1.0015	0.7622
IL-2	0.0180	5.151×e <sup>-5</sup> to 0.5882	0.0164
IL-4	0.0121	2.495×e <sup>-6</sup> to 3.698	0.1477
IL-5	0.4017	0.0787 to 1.1196	0.0873
IL-6	1.0282	0.5270 to 1.8497	0.9261
IL-7	0.1333	0.0133 to 0.6706	0.0089
IL-8	0.9962	0.9652 to 1.0098	0.6278
IL-9	0.1535	0.0070 to 0.8868	0.0180
IL-10	0.9250	0.3274 to 2.2574	0.8672
IL-12	0.0067	3.545×e <sup>-6</sup> to 0.7440	0.0326
IL-13	7.762×e <sup>-5</sup>	4.94×e <sup>-10</sup> to 0.1873	0.0092
IL-17	0.4369	0.0572 to 2.2218	0.3331
Eotaxin/CCL11	0.9626	0.8830 to 1.0327	0.3008
b-FGF	0.5378	0.2200 to 0.9869	0.0444
G-CSF	0.9968	0.9590 to 1.0276	0.8467
GM-CSF	1.128×e <sup>-7</sup>	1.91×e <sup>-21</sup> to 0.1393	0.0141
IFN $\gamma$	0.7547	0.2793 to 1.4865	0.4491
IP-10/CXCL10	0.9976	0.9943 to 0.9997	0.0166
MCP-1/CCL2	0.9773	0.8973 to 1.0293	0.4307
MIP-1 $\alpha$ /CCL3	1.0107	0.8953 to 1.1226	0.8424
PDGF-BB	0.9764	0.9483 to 0.9968	0.0184
MIP-1 $\beta$ /CCL4	0.9859	0.9564 to 1.0068	0.2011
RANTES/CCL5	0.9979	0.9950 to 0.9996	0.0020
TNF- $\alpha$	0.6796	0.3111 to 1.0278	0.0744
Multivariate analysis†			
IL-7	0.2270	0.0177 to 1.5548	0.1384
RANTES/CCL5	0.9980	0.9953 to 0.9998	0.0263
Adjusted by clinical parameters‡			
RANTES/CCL5	0.9970	0.9934 to 0.9993	0.0018
HRCT pattern, diffuse/non-diffuse	19.2129	1.9938 to 438.0654	0.0084

\*OR >1 means an increase in each continuous parameter (pg/mL) indicating high risk of 90-day mortality.

†Multivariate logistic regression analysis with stepwise selection was performed using 10 significant cytokines by univariate analysis with a p value <0.05: IL-2, IL-7, IL-9, IL-12, IL13, b-FGF, GM-CSF, IP-10, PDGF and RANTES.

‡Multivariate logistic regression analysis with stepwise selection was performed using RANTES and HRCT pattern at the time of diagnosis AE and use of antifibrotic drugs before AE.

AE, acute exacerbation; b-FGF, basic fibroblast growth factor; CCL, CC chemokine ligand; CXCL, CXC chemokine ligand; G-CSF, granulocyte colony-stimulating factor; GM-CSF, granulocyte-macrophage colony-stimulating factor; HRCT, high-resolution CT; IFN, interferon; IIPs, idiopathic interstitial pneumonias; IL, interleukin; IP-10, IFN- $\gamma$  inducible protein; MCP, monocyte chemoattractant protein; MIP, macrophage inflammatory protein; PDGF, platelet-derived growth factor; RANTES, regulated on activation, normal T cell expressed and secreted; TNF, tumour necrosis factor.

In this study, we demonstrated that a greater increase in serum IL-10 1 week after the time of diagnosis of AE was a significant predictor of a poor prognosis. IL-10 is a well-known anti-inflammatory cytokine that inhibits production of proinflammatory cytokines and chemokines by monocytes and macrophages.<sup>45</sup> The profibrotic role of IL-10 is controversial. Silica-induced pulmonary fibrosis

was markedly less severe in IL-10 deficient mice than in wild-type mice because of a reduction in lung levels of TGF- $\beta$ , which is a potent inducer of pulmonary fibrosis.<sup>45</sup> However, transient expression of IL-10 by gene transfer for less than 1 week reduced the severity of bleomycin-induced pulmonary fibrosis by reducing production and activation of TGF- $\beta$  in bronchioalveolar lavage fluid

**Table 5** Increase in serum cytokines in patients with AE-IIPs who were alive at 90 days and those who were not\*

Parameters	Survivors (n=16)	Non-survivors (n=6)	P value
$\Delta$ IL-ra	-53.0175 (-297.7100 to -4.1050)	-132.3275 (-371.7150 to 32.3925)	0.8537
$\Delta$ IL-2	0.0050 (-0.0850 to 0.2150)	0.2600 (0.0925 to 0.3300)	0.0356
$\Delta$ IL-4	0.1050 (0.0625 to 0.1875)	0.0600 (-0.0300 to 0.1800)	0.4835
$\Delta$ IL-5	-0.6200 (-1.1050 to 0.0425)	0.5700 (-0.0550 to 1.0225)	0.0183
$\Delta$ IL-6	-0.1950 (-0.3725 to 0.0625)	0.1175 (-0.5975 to 0.6150)	0.3379
$\Delta$ IL-7	-0.3200 (-0.7425 to 0.0675)	0.1100 (-0.2525 to 0.7400)	0.0389
$\Delta$ IL-8	-0.4350 (-2.7925 to 0.9325)	0.5400 (-0.4375 to 6.9350)	0.3020
$\Delta$ IL-9	-0.4300 (-0.9275 to -0.0200)	0.0700 (-0.1800 to 0.5450)	0.0296
$\Delta$ IL-10	-0.5875 (-0.9575 to -0.2275)	0.2200 (0.0100 to 1.3975)	0.0015
$\Delta$ IL-12	-0.0100 (-0.1525 to 0.1400)	0.1200 (-0.0250 to 0.2125)	0.2530
$\Delta$ IL-13	0.1000 (0.0025 to 0.4475)	0.0950 (-0.0150 to 0.2750)	0.7399
$\Delta$ IL-17	0.0450 (-0.1725 to 0.3575)	0.0200 (-0.3625 to 0.6225)	0.7963
$\Delta$ Eotaxin/CCL11	17.7050 (13.0075 to 37.4450)	10.3600 (-4.7950 to 26.0525)	0.1048
$\Delta$ b-FGF	0.3450 (0.1500 to 1.4000)	1.2900 (0.2025 to 3.0475)	0.2381
$\Delta$ G-CSF	-0.0350 (-12.2625 to 1.4288)	2.6325 (0.7013 to 3.1888)	0.0325
$\Delta$ GM-CSF	-0.0175 (-0.0800 to 0.0775)	0.0075 (-0.0150 to 0.0850)	0.1798
$\Delta$ IFN $\gamma$	-0.5350 (-1.7175 to -0.2850)	-0.0900 (-0.2525 to 0.2575)	0.0183
$\Delta$ IP-10/CXCL10	-214.3300 (-570.4550 to 32.1200)	-23.8450 (-393.8175 to 22.3575)	0.4610
$\Delta$ MCP-1/CCL2	-1.5100 (-9.5725 to -0.3175)	5.5650 (2.3700 to 27.1600)	0.0032
$\Delta$ MIP-1 $\alpha$ /CCL3	-0.2050 (-0.3600 to -0.0025)	0.0400 (-0.0700 to 0.3200)	0.0900
$\Delta$ PDGF-BB	-11.0900 (-52.3975 to 12.8250)	-5.800 (-35.8575 to 44.4400)	0.6583
$\Delta$ MIP-1 $\beta$ /CCL4	-109.4150 (-167.2625 to -58.6475)	-88.8850 (-161.6425 to -58.8800)	0.7124
$\Delta$ RANTES/CCL5	-837.7300 (-2753.1075 to -447.1125)	-194.8350 (-1785.9525 to 243.8575)	0.1845
$\Delta$ TNF- $\alpha$	-1.200 (-2.3025 to -0.5375)	-0.4850 (-1.6625 to 0.6525)	0.1404

\*Increase in serum cytokines on day 8 (pg/mL) was compared between survivors and non-survivors on day 91 by the Wilcoxon rank-sum test.

AE, acute exacerbation; b-FGF, basic fibroblast growth factor; CCL, CC chemokine ligand; CXCL, CXC chemokine ligand; G-CSF, granulocyte colony-stimulating factor; GM-CSF, granulocyte-macrophage colony-stimulating factor; IFN, interferon; IIPs, idiopathic interstitial pneumonias; IL, interleukin; IP-10, IFN- $\gamma$  inducible protein; MCP, monocyte chemotactic protein; MIP, macrophage inflammatory protein; PDGF, platelet-derived growth factor; RANTES, regulated on activation, normal T cell expressed and secreted; TNF, tumour necrosis factor.

when compared with mice that underwent control gene transfer.<sup>46 47</sup> These inconsistent reports concerning the effect of IL-10 on fibrosis might reflect a difference in duration of IL-10 expression.

Long-term, lung-specific overexpression of IL-10 in mature mice for 1 and 3 months resulted in infiltration of mononuclear cells into the lung and increased expression of MCP-1/CCL2 mRNA<sup>48</sup>; moreover, pulmonary fibrosis in IL-10 transgenic mice was found to be inhibited by anti-MCP1/CCL2 antibody.<sup>48</sup> Induction of MCP-1/CCL2 by long-term elevation of serum IL-10 in transgenic mice may be consistent with the significant predictive value of  $\Delta$ MCP-1 with regard to 90-day survival noted in our present study. Association between persistent elevation of IL-10 and pulmonary fibrosis in the transgenic mice<sup>48</sup> might explain our finding of a poor prognosis in patients with a marked increase in serum IL-10 levels by day 8. This hypothesis is consistent with the poor survival documented in patients with interstitial lung disease

associated with clinically amyopathic dermatomyositis who are antimelanoma differentiation-associated gene 5 antibody positive and show an increase in serum IL-10 during their clinical course.<sup>49</sup>

The results of this study could alter the treatment strategy for AE-IIPs. Nintedanib<sup>50</sup> is known to suppress the effects of profibrotic cytokines, including VEGF,<sup>51</sup> b-FGF<sup>52</sup> and PDGF.<sup>25</sup> The ability of nintedanib to inhibit AE of IPF was demonstrated in the INPULSYS trial<sup>50</sup>; however, there was no improvement in survival after the time of diagnosis of AE in nintedanib-treated cases.<sup>53</sup> If our findings are correct, use of nintedanib soon after the time of diagnosis of AE might result in a poor survival outcome in patients with AE-IIPs. Higher serum levels of profibrotic cytokines, including b-FGF and PDGF, at the time of diagnosis of AE were significantly good prognostic factors in our patients with AE-IIPs. Hence, administration of nintedanib for more than 7 days after the time of diagnosis of AE rather than early administration



**Table 6** Prognostic significance of an increase in serum cytokine levels at day 8 from baseline levels for AE-IIPs: logistic regression analysis of 90-day mortality

Parameters	OR*	95% CI	P value
Univariate analysis			
ΔIL-ra	1.0003	0.9987 to 1.0038	0.7276
ΔIL-2	6.5971	0.2441 to 453.9501	0.2575
ΔIL-4	0.1854	0.000223 to 68.6763	0.5734
ΔIL-5	1.1894	0.7046 to 2.1647	0.4765
ΔIL-6	2.1743	0.8055 to 14.1524	0.1614
ΔIL-7	12.6586	1.5148 to 407.2435	0.0130
ΔIL-8	1.0164	0.9928 to 1.0842	0.3159
ΔIL-9	3.7582	0.9318 to 34.8311	0.0692
ΔIL-10	489.7742	5.4452 to 3 967 492	0.0002
ΔIL-12	0.9146	0.0764 to 4.1373	0.9122
ΔIL-13	0.3314	0.0034 to 4.5552	0.4703
ΔIL-17	0.7084	0.0691 to 4.7883	0.7327
ΔEotaxin/CCL11	0.9306	0.8249 to 1.0037	0.0657
Δb-FGF	1.4213	0.7410 to 2.9857	0.2883
ΔG-CSF	1.2878	1.0029 to 2.1824	0.0418
ΔGM-CSF	2.6346	0.0085 to 996.8476	0.7260
ΔIFN $\gamma$	24.2498	1.7469 to 4292.726	0.0063
ΔIP-10/CXCL10	1.0002	0.9986 to 1.0018	0.7270
ΔMCP-1/CCL2	1.4248	1.0812 to 2.2406	0.0009
ΔMIP-1 $\alpha$ /CCL3	2.4528	0.9378 to 28.5121	0.1541
ΔPDGF-BB	1.0104	0.9931 to 1.0346	0.2433
ΔMIP-1 $\beta$ /CCL4	1.0048	0.9970 to 1.0211	0.2765
ΔRANTES/CCL5	1.0001	0.9999 to 1.0007	0.2813
ΔTNF- $\alpha$	2.1615	0.9022 to 7.3186	0.0918
Multivariate analysis†			
ΔIL-10	489.7	5.445 to 3 967 492	0.0002

\*OR >1 means an increase in each parameter (pg/mL) indicating a high risk of 90-day mortality.

†Multivariate logistic regression analysis with stepwise selection was performed using five significant ( $p < 0.05$ ) cytokines by univariate analysis: ΔIL-7, ΔIL10, ΔG-CSF, ΔIFN- $\gamma$  and ΔMCP-1.

b-FGF, basic fibroblast growth factor; CCL, CC chemokine ligand; CXCL, CXC chemokine ligand; G-CSF, granulocyte colony-stimulating factor; GM-CSF, granulocyte-macrophage colony-stimulating factor; IFN, interferon; IIPs, idiopathic interstitial pneumonias; IL, interleukin; IP-10, IFN- $\gamma$  inducible protein; MCP, monocyte chemotactic protein; MIP, macrophage inflammatory protein; PDGF, platelet-derived growth factor; RANTES, regulated on activation, normal T cell expressed and secreted; TNF, tumour necrosis factor.

might confer a survival advantage in AE-IPF. In addition, we have shown AE-IIPs with use of antifibrotic drugs before AE showed worse survival than those without their use (table 2 and online supplemental table 3). This can be explained by difference in severity of patients of IIPs before AE with or without use of antifibrotic drugs in our study (online supplemental table 3); however, cessation of antifibrotic drugs temporally during the 7 days after the AE diagnosis might be better to improve prognosis of AE-IIPs from our hypothesis. How to use antifibrotic drugs for AE-IPF patients is important problem to be solved by future studies.

Effects of thrombomodulin on cytokine production have not been clarified sufficiently; however, they have

been investigated for other diseases except for AE-IIPs. Kudo *et al*<sup>54</sup> reported IL-10 and TGF- $\beta$  in the lung homogenate of lipopolysaccharide induced acute respiratory distress syndrome (ARDS) model mice treated by rhTM was significantly elevated 24 hours after the onset of ARDS as compared with control mice without rhTM treatment. Temporal elevation of these two cytokines might help recovery from the epithelial injury and inflammation at AE as we have shown from our data. In addition, rhTM might inhibit endothelial cell injury and help reduction of RANTES/CCL5 as shown in patients treated with haematopoietic stem cell transplantation.<sup>55 56</sup> When and how long to administer rhTM might be important to improve the prognosis of AE-IIP.



This study has several limitations. First, the numbers of patients with AE-IPF and AE-non-IPF were too small to draw definitive conclusion. We cannot divide subjects into original and validation cohorts. We cannot separately analyse AE-IPF and AE-non-IPF to reduce bias, either. Second, serum samples were obtained from patients with AE-IIPs treated with rhTM and corticosteroids of SETUP trial; however, serum samples could not be obtained from all the subjects of the trial. Third, a randomised controlled trial of rhTM could not clarify its efficacy in AE-IPF,<sup>57</sup> and we cannot recommend it as a treatment for AE-IPF or AE-IIPs based on the present evidence. Therefore, our present findings need to be confirmed in a study of patients with AE-IIPs who are treated with corticosteroids only. Fourth, the levels of several cytokines could not be measured simultaneously using the Bio-Plex system. Fifth, PMX/DHP therapy is known to reduce various serum cytokine levels<sup>13</sup> and might have affected the serum cytokine profile on day 8; however, only one case was treated with the therapy.

## CONCLUSIONS

An elevated serum RANTES/CCL5 level at the time of diagnosis of AE predicted a good survival outcome in patients with AE-IIPs and a marked increase in serum IL-10 on day 8 predicted a poor survival outcome. Therefore, early elevation of serum RANTES/CCL5 and a rapid reduction of IL-10 could predict 90-day survival in patients with AE-IIPs treated with corticosteroids and rhTM.

## Author affiliations

<sup>1</sup>Clinical Research Center, National Hospital Organization Kinki-Chuo Chest Medical Center, Sakai City, Japan

<sup>2</sup>Department of Respiratory Medicine, Osaka Prefectural Hospital Organization Osaka Habikino Medical Center, Habikino City, Japan

<sup>3</sup>Department of Respiratory Medicine, National Hospital Organization Osaka Toneyama Medical Center, Toyonaka City, Japan

<sup>4</sup>Department of Respiratory Medicine and Clinical Immunology, Osaka University Graduate School of Medicine, Suita City, Japan

<sup>5</sup>Department of Respiratory Medicine, National Hospital Organization Osaka Minami Medical Center, Kawachinagano City, Japan

<sup>6</sup>Department of Critical Care Medicine, Yao Tokushukai Hospital, Yao City, Japan

**Acknowledgements** We are grateful to Ms Homma T, Ms Hirakawa C and Ms Matsui Y for their secretarial work and to Ms Imai Y for the data management work. We would like to thank Editage ([www.editage.com](http://www.editage.com)) for English language editing.

**Contributors** TA contributed to study conception and design, radiological findings, data analysis and interpretation and manuscript preparation. TA had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. HM, SY, YO, CS, KT and HK contributed to study conception and design, data acquisition, data analysis and interpretation and manuscript preparation. YI and MM contributed to study conception and design, data analysis and interpretation and manuscript preparation. MH and KH contributed to data acquisition, data analysis and interpretation and manuscript preparation. MA contributed to the radiological findings and manuscript preparation. All authors approved the final version of the manuscript.

**Funding** This study was partially supported by a JSPS KAKENHI Grant Number JP17K09636 awarded to TA, YI and MH, a National Hospital Organization (H28-NHO (Kokyu)-2) grant awarded to TA and YI, and AMED grants (DLD/14526278 and PAP/14526182) awarded to YI and to YI and TA, respectively.

**Competing interests** HM, MH, HK, SY, YO, MM, KH, CS, KT and MA have no competing interests. YI is an advisor to Boehringer Ingelheim, Shionogi and Asahi Kasei. TA has received lecture fees from Boehringer Ingelheim and Shionogi for activities not connected with the submitted work.

**Patient and public involvement statement** This research was done without patient involvement. Patients were not invited to comment on the study design and were not consulted to develop patient relevant outcomes or interpret the results. Patients were not invited to contribute to the writing or editing of this document for readability or accuracy.

**Patient consent for publication** Not required.

**Ethics approval** This study was approved by the Institutional Review Board of the Kinki-Chuo Chest Medical Center (KCCMC; approval number: 447, 24 February 2014), as well as the review boards of the other institutions. Written informed consent was obtained from all participants in the rhTM arm.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** No data are available. The data that support the findings of this study have not been made available because this study was approved in 2014 and data sharing was not included in the design of this study.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Toru Arai <http://orcid.org/0000-0002-2222-0683>

Yoshikazu Inoue <http://orcid.org/0000-0003-3994-874X>

## REFERENCES

- Raghu G, Collard HR, Egan JJ. On behalf of the ATS/ERS/JRS/ALAT Committee on idiopathic pulmonary fibrosis. An official ATS/ERS/JRS/ALAT statement: idiopathic pulmonary fibrosis: evidence-based guidelines for diagnosis and management. *Am J Respir Crit Care Med* 2011;183:788–824.
- Raghu G, Remy-Jardin M, Myers JL, et al. Diagnosis of idiopathic pulmonary fibrosis. An official ATS/ERS/JRS/ALAT clinical practice guideline. *Am J Respir Crit Care Med* 2018;198:e44–68.
- Akira M, Kozuka T, Yamamoto S, et al. Computed tomography findings in acute exacerbation of idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2008;178:372–8.
- Collard HR, Ryerson CJ, Corte TJ, et al. Acute exacerbation of idiopathic pulmonary fibrosis. An international Working Group report. *Am J Respir Crit Care Med* 2016;194:265–75.
- Arai T, Kagawa T, Sasaki Y, et al. Heterogeneity of incidence and outcome of acute exacerbation in idiopathic interstitial pneumonia. *Respirology* 2016;21:1431–7.
- Suzuki A, Kondoh Y, Brown KK, et al. Acute exacerbations of fibrotic interstitial lung diseases. *Respirology* 2020;25:525–34.
- Saito H, Maruyama I, Shimazaki S, et al. Efficacy and safety of recombinant human soluble thrombomodulin (ART-123) in disseminated intravascular coagulation: results of a phase III, randomized, double-blind clinical trial. *J Thromb Haemost* 2007;5:31–41.
- Kataoka K, Taniguchi H, Kondoh Y, et al. Recombinant human thrombomodulin in acute exacerbation of idiopathic pulmonary fibrosis. *Chest* 2015;148:436–433.
- Arai T, Kida H, Ogata Y, et al. Recombinant thrombomodulin for acute exacerbation in idiopathic interstitial pneumonias. *Respirology* 2019;24:658–66.
- Okamoto T, Tanigami H, Suzuki K, et al. Thrombomodulin: a bifunctional modulator of inflammation and coagulation in sepsis. Hindawi publishing Corporation critical care research and practice 2012, article ID 61545, 10 Pages.

- 11 Collard HR, Calfee CS, Wolters PJ, *et al.* Plasma biomarker profiles in acute exacerbation of idiopathic pulmonary fibrosis. *Am J Physiol Lung Cell Mol Physiol* 2010;299:L3–7.
- 12 Konishi K, Gibson KF, Lindell KO. Gene expression profile of acute exacerbations of idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2009;189:167–75.
- 13 Oishi K, Mimura-Kimura Y, Miyasho T, *et al.* Association between cytokine removal by polymyxin B hemoperfusion and improved pulmonary oxygenation in patients with acute exacerbation of idiopathic pulmonary fibrosis. *Cytokine* 2013;61:84–9.
- 14 Papisir SA, Tomos IP, Karakatsani A, *et al.* High levels of IL-6 and IL-8 characterize early-on idiopathic pulmonary fibrosis acute exacerbations. *Cytokine* 2018;102:168–72.
- 15 Tachibana K, Inoue Y, Nishiyama A, *et al.* Polymyxin-B hemoperfusion for acute exacerbation of idiopathic pulmonary fibrosis: serum IL-7 as a prognostic marker. *Sarcoidosis Vasc Diffuse Lung Dis* 2011;28:113–22.
- 16 Taniguchi H, Ebina M, Kondoh Y, *et al.* Pirfenidone in idiopathic pulmonary fibrosis. *Eur Respir J* 2010;35:821–9.
- 17 Travis WD, Costabel U, Hansell DM, *et al.* An official American thoracic Society/European respiratory Society statement: update of the International multidisciplinary classification of the idiopathic interstitial pneumonias. *Am J Respir Crit Care Med* 2013;188:733–48.
- 18 Celli BR, MacNee W, ATS/ERS Task Force. Standards for the diagnosis and treatment of patients with COPD: a summary of the ATS/ERS position paper. *Eur Respir J* 2004;23:932–46.
- 19 Gando S, Wada H, Asakura H, *et al.* Evaluation of new Japanese diagnostic criteria for disseminated intravascular coagulation in critically ill patients. *Clin Appl Thromb Hemost* 2005;11:71–6.
- 20 Bernard GR, Artigas A, Brigham KL, *et al.* The American-European consensus conference on ARDS: definitions, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med* 1994;149:818–24.
- 21 Wang JH, Devalia JL, Xia C, Raymond CX, *et al.* Expression of RANTES by human bronchial epithelial cells in vitro and in vivo and the effect of corticosteroids. *Am J Respir Cell Mol Biol* 1996;14:27–35.
- 22 Yoshioka S, Mukae H, Sugiyama K, *et al.* High-BAL fluid concentrations of RANTES in nonspecific interstitial pneumonia compared with usual interstitial pneumonia. *Respir Med* 2004;98:945–51.
- 23 Kodama N, Yamaguchi E, Hizawa N, *et al.* Expression of RANTES by bronchoalveolar lavage cells in nonsmoking patients with interstitial lung diseases. *Am J Respir Cell Mol Biol* 1998;18:526–31.
- 24 Pierce EM, Carpenter K, Jakubczik C, *et al.* Idiopathic pulmonary fibrosis fibroblasts migrate and proliferate to CC chemokine ligand 21. *Eur Respir J* 2007;29:1082–93.
- 25 Sahin H, Wasmuth HE. Chemokines in tissue fibrosis. *Biochim Biophys Acta* 2013;1832:1041–8.
- 26 Nishioka Y, Azuma M, Kishi M, *et al.* Targeting platelet-derived growth factor as a therapeutic approach in pulmonary fibrosis. *J Med Invest* 2013;60:175–83.
- 27 Chung SI, Horton JA, Ramalingam TR, *et al.* IL-13 is a therapeutic target in radiation lung injury. *Sci Rep* 2016;6:39714.
- 28 Huang M, Sharma S, Zhu LX, *et al.* IL-7 inhibits fibroblast TGF-beta production and signaling in pulmonary fibrosis. *J Clin Invest* 2002;109:931–7.
- 29 Zhang L, Keane MP, Zhu LX, *et al.* Interleukin-7 and transforming growth factor-beta play counter-regulatory roles in protein kinase C-delta-dependent control of fibroblast collagen synthesis in pulmonary fibrosis. *J Biol Chem* 2004;279:28315–9.
- 30 Moore BB, Coffey MJ, Christensen P. Gm-CSF regulates bleomycin-induced pulmonary fibrosis via a prostaglandin-dependent mechanism. *J Immunology* 2000;165:4032–9.
- 31 Tager AM, Kradin RL, LaCamera P, *et al.* Inhibition of pulmonary fibrosis by the chemokine IP-10/CXCL10. *Am J Respir Cell Mol Biol* 2004;31:395–404.
- 32 van den Brùle S, Heymans J, Havaux X, *et al.* Profibrotic effect of IL-9 overexpression in a model of airway remodeling. *Am J Respir Cell Mol Biol* 2007;37:202–9.
- 33 Arras M, Huaux F, Vink A, *et al.* Interleukin-9 reduces lung fibrosis and type 2 immune polarization induced by silica particles in a murine model. *Am J Respir Cell Mol Biol* 2001;24:368–75.
- 34 Geiser T. Idiopathic pulmonary fibrosis—a disorder of alveolar wound repair? *Swiss Med Wkly* 2003;133:405–11.
- 35 Wijsenbeek M, Cottin V. Spectrum of fibrotic lung diseases. *N Engl J Med* 2020;383:958–68.
- 36 Yano T, Mason RJ, Pan T, *et al.* Kgf regulates pulmonary epithelial proliferation and surfactant protein gene expression in adult rat lung. *Am J Physiol Lung Cell Mol Physiol* 2000;279:L1146–58.
- 37 Sugiyama M, Kinoshita N, Isa S. Serum CCL17 level become a predictive marker to distinguish between mild/moderate and severe/critical disease in patients with COVID-19. *Gene*;201:145145.
- 38 Belperio JA, Dy M, Murray L, *et al.* The role of the Th2 CC chemokine ligand CCL17 in pulmonary fibrosis. *J Immunol* 2004;173:4692–8.
- 39 Yogo Y, Fujishima S, Inoue T, *et al.* Macrophage derived chemokine (CCL22), thymus and activation-regulated chemokine (CCL17), and CCR4 in idiopathic pulmonary fibrosis. *Respir Res* 2009;10:80.
- 40 Xie L, Zhao L. Role of CXCL12/CXCR-4-mediated circulating fibrocytes in pulmonary fibrosis. *J Biomedicine* 2017;2:134–9.
- 41 Moeller A, Gilpin SE, Ask K, *et al.* Circulating fibrocytes are an indicator of poor prognosis in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2009;179:588–94.
- 42 Strieter RM, Keeley EC, Hughes MA, *et al.* The role of circulating mesenchymal progenitor cells (fibrocytes) in the pathogenesis of pulmonary fibrosis. *J Leukoc Biol* 2009;86:1111–8.
- 43 Isgrò M, Bianchetti L, Marini MA, *et al.* The C-C motif chemokine ligands CCL5, CCL11, and CCL24 induce the migration of circulating fibrocytes from patients with severe asthma. *Mucosal Immunol* 2013;6:718–27.
- 44 Tai W, Xu Y, Ding J, *et al.* Fibrocytes ameliorate acute lung injury by decreasing inflammatory cytokine and chemokine levels and reducing neutrophil accumulation in the lung. *Cell Physiol Biochem* 2017;44:1526–36.
- 45 Barbarin V, Arras M, Misson P, *et al.* Characterization of the effect of interleukin-10 on silica-induced lung fibrosis in mice. *Am J Respir Cell Mol Biol* 2004;31:78–85.
- 46 Nakagome K, Dohi M, Okunishi K, *et al.* In vivo IL-10 gene delivery attenuates bleomycin induced pulmonary fibrosis by inhibiting the production and activation of TGF-beta in the lung. *Thorax* 2006;61:886–94.
- 47 Arai T, Abe K, Matsuoka H, *et al.* Introduction of the interleukin-10 gene into mice inhibited bleomycin-induced lung injury in vivo. *Am J Physiol Lung Cell Mol Physiol* 2000;278:L914–22.
- 48 Spight D, Zho B, Haas M. Immunoregulatory effects of regulated, lung-target expression of IL-10 in vivo. *Am J Physiol Cell Mol Physiol* 2005;288:L251–65.
- 49 Takada T, Ohashi K, Hayashi M, *et al.* Role of IL-15 in interstitial lung diseases in amyopathic dermatomyositis with anti-MDA-5 antibody. *Respir Med* 2018;141:7–13.
- 50 Richeldi L, du Bois RM, Raghu G, *et al.* Efficacy and safety of nintedanib in idiopathic pulmonary fibrosis. *N Engl J Med* 2014;370:2071–81.
- 51 Hamada N, Kuwano K, Yamada M, *et al.* Anti-Vascular endothelial growth factor gene therapy attenuates lung injury and fibrosis in mice. *J Immunol* 2005;175:1224–31.
- 52 Inoue Y, King TE, Tinkle SS, *et al.* Human mast cell basic fibroblast growth factor in pulmonary fibrotic disorders. *Am J Pathol* 1996;149:2037–54.
- 53 Collard HR, Richeldi L, Kim DS, *et al.* Acute exacerbations in the INPULSIS trials of nintedanib in idiopathic pulmonary fibrosis. *Eur Respir J* 2017;49:1601339.
- 54 Kudo D, Toyama M, Aoyagi T, *et al.* Involvement of high mobility group box 1 and the therapeutic effect of recombinant thrombomodulin in a mouse model of severe acute respiratory distress syndrome. *Clin Exp Immunol* 2013;173:276–87.
- 55 Nomura S, Ishii K, Fujita S, *et al.* Associations between acute GVHD-related biomarkers and endothelial cell activation after allogeneic hematopoietic stem cell transplantation. *Transpl Immunol* 2017;43:44:27–32.
- 56 Ishii K, Shimizu M, Nomura S. Prophylaxis against endothelial damage using recombinant human soluble thrombomodulin (rTM) in allogeneic hematopoietic stem cell transplantation (allo-HSCT). *Blood* 2010;116:4523.
- 57 Kondoh Y, Azuma A, Inoue Y, *et al.* Thrombomodulin alfa for acute exacerbation of idiopathic pulmonary fibrosis. A randomized, double-blind placebo-controlled trial. *Am J Respir Crit Care Med* 2020;201:1110–9.

**Online supplement**

**Prognostic Significance of Serum Cytokines during Acute Exacerbation of  
Idiopathic Interstitial Pneumonias treated with Thrombomodulin**



**Online supplemental Table S1.** Patient characteristics of the subjects with serum sample in this study compared with AE-IIPs without serum sample in SETUP trial\*

Parameters	Sample obtained (n=28)	Sample not obtained (n=11)	p-value
<b>Stable state</b>			
IPF/Non-IPF	8/20	4/7	0.7086
Sex, Male/Female	21/7	7/4	0.6940
Smoking, y/n	19/9	8/3	1.0000
mMRC, $\leq 1 / \geq 2$	10/18	5/6	0.7181
Stage, I - III / IV	17/11	7/4	1.0000
LTOT before AE, y/n	11/17	3/8	0.7129
Prednisolone before AE, y/n	9/19	6/5	0.2773
Antifibrotic drugs, y/n	5 <sup>#</sup> /23	1 <sup>§</sup> /10	0.6548
<b>At the time of AE diagnosis</b>			
Age, yrs	74.5 (69.75–79)	72 (64–80)	0.2170
HRCT pattern, diffuse/non- diffuse	12/16	3/8	0.4770
PaO <sub>2</sub> /FiO <sub>2</sub> ratio, $\leq 200 / > 200$	17/11	6/5	0.7338
WBC, / $\mu$ L	10,150 (9,050–12,950)	10,700 (8,800–14,340)	0.9378
LDH, U/L	317.0 (245.5–404.5)	381.0 (290.0–451.0)	0.3030
KL-6, U/mL	1,196 (889–2,142)	859 (616–2,251)	0.2679
CRP, mg/dL	12.38 (5.07–14.28)	12.36 (5.05–14.24)	1.0000
FDP, $< 10 / \geq 10$ mg/L	21/7	7/4	0.6940
90-day survival, y/n	19/9	7/4	1.0000

AE, acute exacerbation; CRP, C-reactive protein; FDP, fibrin degradation product; FiO<sub>2</sub>, fraction of inspired oxygen; HRCT, high-resolution computed tomography; IPF, idiopathic pulmonary fibrosis; IIPs, idiopathic interstitial pneumonias; KL-6, Krebs von den Lungen-6; LDH, lactate dehydrogenase; LTOT, long-term oxygen therapy; mMRC, modified Medical Research Council score; PaO<sub>2</sub>, arterial oxygen tension.

\*Each parameter was compared between the two groups using Fisher's exact test or the Wilcoxon rank-sum test.

#: Pirfenidone (n=5), nintedanib (n=1). In one case, pirfenidone was firstly prescribed and nintedanib was secondary administered. §: Pirfenidone (n=1).

**Online supplementary Table S2.** Therapy for AE-IIPs and management of respiratory failure

Therapy and Management	N=28
<b>Therapy</b>	
Steroid pulse, y/n	25/3
Prednisolone, y/n*	27/1
Dose of prednisolone**, mg/kg/day	0.980 (0.909–1.017)
Immunosuppressant, y/n	5/23
Pirfenidone, y/n	3/25
Nintedanib, y/n	0/28
Antifibrotic drugs#, y/n	3/25
Empiric antibiotics, y/n	28/0
Macrolide, y/n	5/23
Antacids, y/n	27/1
PMX-DHP therapy, y/n	1/27
Thrombomodulin, y/n	28/0
Anticoagulants#, y/n	6/22
Anti-platelet drugs#, y/n	5/23
<b>Management of respiratory failure</b>	
IPPV, y/n	4/24
NPPV, y/n	11/17
IPPV or NPPV, y/n	13/15
NHF, y/n	9/19
NHF without IPPV, NPPV, y/n	5/23
IPPV or NPPV or NHF, y/n	18/10

Abbreviations: AE, acute exacerbation; IIPs, idiopathic interstitial pneumonias; rhTM, recombinant human soluble thrombomodulin; AZP, azathioprine; CyA, cyclosporine A; CPA, cyclophosphamide; IPPV, invasive positive pressure ventilation; NPPV, non-invasive positive pressure ventilation; NHF, nasal high flow therapy.

\*AE-IIPs patients with deceased and with no administration of prednisolone during a high-dose methylprednisolone therapy.

\*\*Prednisolone dose is described as the median (IQR).

#: Anticoagulants and anti-platelet drugs were used for comorbidities.

**Online supplementary Table S3.** Clinical parameters according to the usage of antifibrotic drugs before the diagnosis of AE-IIPs (n=28)\*

Clinical parameters	Usage of antifibrotic drugs (n=5)	No usage of antifibrotic drugs (n=23)	p-value
<b>Stable state</b>			
Gender, Male/Female	4/1	17/6	1.0000
IPF/Non-IPF	2/3	6/17	0.6056
Smoking, y/n	3/2	16/7	1.0000
mMRC, $\leq 1 / \geq 2$	1/4	9/14	0.6264
Stage, I - III / IV	0/5	17/6	0.0047
LTOT before AE, y/n	5/0	6/17	0.0047
Prednisolone before AE, y/n	1/4	8/15	1.0000
<b>At the time of AE diagnosis</b>			
PaO <sub>2</sub> /FiO <sub>2</sub> ratio, $\leq 200 / > 200$	3/2	14/9	1.0000
HRCT, diffuse/non-diffuse	3/2	9/14	0.6239
<b>Outcome</b>			
90-day survival, y/n	1/4	18/5	0.0256

Abbreviations; AE, acute exacerbation; FiO<sub>2</sub>, fraction of inspired oxygen; HRCT, high-resolution computed tomography; IPF, idiopathic pulmonary fibrosis; IIPs, idiopathic interstitial pneumonias; LTOT, long-term oxygen therapy; mMRC, modified Medical Research Council score; PaO<sub>2</sub>, arterial oxygen tension.

\*: Association between usage of antifibrotic drugs and clinical parameters at the stable state and at the time of diagnosis of AE-IIPs was evaluated by Fisher's exact test.

**Online supplementary Table S4.** Serum cytokines of AE-IIPs, Between IPF and non-IPF (n=28)\*

Parameters	AE-IPF (n = 8)	AE-non-IPF (n = 20)	p-value
IL-ra	61.29 (26.06–179.92)	163.69 (8.45–426.18)	0.6274
IL-2	0.405 (0.307–1.072)	0.380 (0.240–0.685)	0.5579
IL-4	0.305 (0.197–0.415)	0.285 (0.197–0.405)	0.9797
IL-5	1.430 (1.010–2.960)	1.265 (0.930–2.097)	0.7794
IL-6	0.815 (0.250–2.132)	0.650 (0.225–1.822)	0.6654
IL-7	1.245 (0.660–2.137)	1.135 (0.815–1.767)	0.8786
IL-8	2.960 (1.895–46.822)	4.055 (3.030–6.877)	0.7993
IL-9	2.010 (1.787–3.077)	1.975 (1.810–2.860)	0.9594
IL-10	0.880 (0.497–2.060)	0.785 (0.537–1.677)	0.7026
IL-12	0.225 (0.120–0.432)	0.140 (0.067–0.390)	0.6441
IL-13	0.200 (0.082–0.447)	0.130 (0.062–0.307)	0.2728
IL-17	1.065 (0.727–1.722)	1.090 (0.870–1.375)	0.9189
Eotaxin/CCL11	16.890 (9.387–25.122)	18.36 (10.93–30.65)	0.6841
b-FGF	6.025 (4.912–7.772)	5.110 (4.720–6.720)	0.4263
G-CSF	1.560 (0.980–59.422)	4.550 (0.690–13.575)	0.6829
GM-CSF	0.050 (0.025–0.257)	0.040 (0.020–0.132)	0.5993
IFN $\gamma$	0.620 (0.400–2.037)	0.820 (0.542–2.047)	0.4006
IP-10/CXCL10	369.79 (154.25–1111.44)	546.2 (236.9–1143.0)	0.5085
MCP-1/CCL2	4.075 (2.360–18.022)	6.375 (3.555–17.785)	0.1860
MIP-1 $\alpha$ /CCL3	0.540 (0.412–17.267)	0.685 (0.352–1.165)	0.5933
PDGF-BB	67.690 (36.397–100.035)	79.295 (49.235–112.792)	0.7219
MIP-1 $\beta$ /CCL4	246.25 (205.50–490.29)	230.64 (187.34–305.07)	0.4765
RANTES/CCL5	1863 (1672–4891)	1859 (1466–3061)	0.7602
TNF- $\alpha$	4,855 (3.990–11.712)	4.385 (3.865–6.170)	0.5931

Abbreviations; AE, acute exacerbation; IIPs, idiopathic interstitial pneumonias; IPF, idiopathic pulmonary fibrosis; IL, interleukin; CCL, CC chemokine ligand; bFGF, basic fibroblast growth factor; G-CSF; granulocyte colony-stimulating factor; GM-CSF, granulocyte-macrophage colony-stimulating factor; IFN, interferon; IP-10, IFN- $\gamma$  inducible protein; CXCL, CXC chemokine ligand; MCP, monocyte chemotactic protein; MIP, macrophage inflammatory protein; PDGF, platelet-derived growth factor; RANTES, regulated on activation, normal T-cell expressed and secreted; TNF, tumor necrosis factor.

\*: Serum cytokine levels of AE-IPF and AE-non-IPF (pg/mL) were compared with Wilcoxon rank sum test.



**Online supplementary Table S5.**

Association of serum RANTES/CCL5 levels and clinical parameters at the diagnosis of AE-IIPs (n=28)\*

Parameters	rho	p-value
<b>Stable state</b>		
Gender, Male/Female	0.0204	0.9178
IPF/Non-IPF	0.0587	0.7666
Smoking, y/n	0.0189	0.9238
mMRC, $\leq 1 / \geq 2$	0.1015	0.6073
Stage, I - III / IV	-0.3622	0.0582
LTOT before AE, y/n	-0.3622	0.0582
Prednisolone before AE, y/n	0.0473	0.8109
Antifibrotic drugs, y/n	-0.5484**	0.0025
<b>At the time of AE diagnosis</b>		
WBC, / $\mu$ L	0.0236	0.9053
CRP, mg/dL	0.2275	0.2444
LDH, U/L	-0.0616	0.7555
KL-6, U/mL	-0.2740	0.1583
PaO <sub>2</sub> /FiO <sub>2</sub> ratio, $\leq 200 / > 200$	0.2128	0.2770
HRCT, diffuse/non-diffuse	-0.1966	0.3160
FDP, $< 10 / \geq 10$ mg/dL	-0.0255	0.8974

Abbreviations: RANTES, regulated on activation, normal T-cell expressed and secreted; AE, acute exacerbation; IIPs, idiopathic interstitial pneumonias; IPF, idiopathic pulmonary fibrosis; mMRC, modified Medical Research Council score; LTOT, long-term oxygen therapy; WBC, white blood cell; CRP, C-reactive protein; LDH, lactate dehydrogenase; KL-6, Krebs von den Lungen-6; PaO<sub>2</sub>, arterial oxygen tension; FiO<sub>2</sub>, fraction of inspired oxygen; HRCT, high-resolution computed tomography; FDP, fibrin degradation product.

\*: Correlation between serum levels of RANTES/CCL5 at the diagnosis of AE (pg/mL) and clinical parameters at AE and stable state was examined by Spearman rank correlation.

\*\* : This result suggested serum levels of RANTES/CCL5 at the diagnosis of AE was significantly lower in patients with AE-IIPs treated with antifibrotic drugs before the diagnosis of AE than the other patients with AE-IIPs.

**Online supplementary Table S6.** Serum cytokines changes from baseline to day 8 (n=22)\*

Parameters	AE-IIPs at baseline	At day 8	p-value
IL-ra	82.81 (11.83–392.45)	41.41 (7.60–77.64)	0.0025
IL-2	0.405 (0.240–0.745)	0.500 (0.347–0.745)	0.0551
IL-4	0.300 (0.212–0.417)	0.410 (0.290–0.552)	0.0010
IL-5	1.265 (0.920–2.215)	1.070 (0.607–1.645)	0.3790
IL-6	0.480 (0.212–1.425)	0.350 (0.265–0.500)	0.2189
IL-7	1.245 (0.860–1.845)	1.100 (0.630–1.710)	0.0530
IL-8	3.410 (2.412–6.397)	3.695 (2.225–6.475)	0.7417
IL-9	2.115 (1.870–3.060)	2.005 (1.650–2.797)	0.0358
IL-10	0.735 (0.492–1.670)	0.480 (0.185–0.820)	0.0284
IL-12	0.225 (0.120–0.405)	0.230 (0.142–0.460)	0.5332
IL-13	0.165 (0.085–0.350)	0.210 (0.115–0.602)	0.0064
IL-17	1.090 (0.870–1.465)	1.240 (0.950–1.702)	0.3334
Eotaxin/CCL11	18.36 (11.40–31.90)	32.91 (22.01–56.75)	<0.001
b-FGF	5.490 (4.720–7.850)	6.530 (5.800–8.577)	0.0029
G-CSF	2.725 (0.690–10.065)	3.685 (2.185–8.550)	0.7299
GM-CSF	0.050 (0.020–0.157)	0.012 (0.005–0.165)	0.7411
IFN $\gamma$	0.755 (0.545–2.002)	0.300 (0.242–0.602)	0.0003
IP-10/CXCL10	546.2 (205.9–1119.7)	281.7 (155.1–615.9)	0.0164
MCP-1/CCL2	5.460 (3.160–16.035)	5.535 (3.790–12.717)	0.8140
MIP-1 $\alpha$ /CCL3	0.615 (0.352–1.035)	0.445 (0.292–1.090)	0.1575
PDGF-BB	85.45 (52.89–114.12)	61.72 (39.51–161.6)	0.1256
MIP-1 $\beta$ /CCL4	233.4 (191.7–301.0)	128.0 (111.6–161.6)	<0.0001
RANTES/CCL5	1883 (1750–3260)	1480 (1233–2058)	0.0010
TNF- $\alpha$	4.385 (3.915–6.345)	3.855 (3.072–4.612)	<0.0001

Abbreviations; AE, acute exacerbation; IIPs, idiopathic interstitial pneumonias; IL, interleukin; CCL, CC chemokine ligand; bFGF, basic fibroblast growth factor; G-CSF; granulocyte colony-stimulating factor; GM-CSF, granulocyte-macrophage colony-stimulating factor; IFN, interferon; IP-10, IFN- $\gamma$  inducible protein; CXCL, CXC chemokine ligand; MCP, monocyte chemotactic protein; MIP, macrophage inflammatory protein; PDGF, platelet-derived growth factor; RANTES, regulated on activation, normal T-cell expressed and secreted; TNF, tumor necrosis factor.

\*: Serum levels of each cytokine at AE-onset (pg/mL) were compared with those on day 8 by paired Wilcoxon rank sum test.

**Online supplementary Table S7.**

Association of serum  $\Delta$ IL-10\* levels and clinical parameters at the onset of AE-IIPs (n=22)\*\*


Parameters	rho	p-value
<b>Stable state</b>		
Gender, Male/Female	-0.0483	0.8311
IPF/Non-IPF	-0.3591	0.1007
Smoking, y/n	-0.1126	0.6117
mMRC, $\leq 1 / \geq 2$	-0.0805	0.7219
Stage, I -III / IV	0.0223	0.9214
LTOT before AE, y/n	0.0223	0.9214
Prednisolone before AE	0.0154	0.9458
Antifibrotic drugs, y/n	-0.3241	0.1412
<b>At the time of AE diagnosis</b>		
WBC, / $\mu$ L	-0.0385	0.8650
CRP, mg/dL	-0.1226	0.5869
LDH, U/L	0.2576	0.2471
KL-6, U/mL	0.2299	0.3034
PaO <sub>2</sub> /FiO <sub>2</sub> ratio, $\leq 200 / > 200$	-0.0573	0.7999
HRCT, diffuse/non-diffuse	0.1341	0.5519
FDP, $< 10 / \geq 10$ mg/dL	0.2309	0.3013

Abbreviations: AE, acute exacerbation; IIPs, idiopathic interstitial pneumonias; RANTES, regulated on activation, normal T-cell expressed and secreted; AE, acute exacerbation; IIP, idiopathic interstitial pneumonia; IPF, idiopathic pulmonary fibrosis; mMRC, modified Medical Research Council score; LTOT, long-term oxygen therapy; WBC, white blood cell; CRP, C-reactive protein; LDH, lactate dehydrogenase; KL-6, Krebs von den Lungen-6; PaO<sub>2</sub>, arterial oxygen tension; FiO<sub>2</sub>, fraction of inspired oxygen; HRCT, high-resolution computed tomography; FDP, fibrin degradation product.

\*:  $\Delta$ IL-10 means increase in serum levels of IL-10 on day 8 compared with those at AE onset.

\*\* : Correlation between serum levels of  $\Delta$ IL-10 on day 8 (pg/mL) and clinical parameters at AE and stable state was examined by Spearman rank correlation.

## Online supplementary Figure S1



Subjects of this study were selected from those in the SETUP trial. From October 2014 to March 2016, 40 cases of AE-IIPs were prospectively enrolled and treated with rhTM and conventional therapy. One case was excluded from the trial, with 39 cases included in the rhTM arm. As the control arm, 67 consecutive AE-IIPs cases were diagnosed and treated with conventional therapy without rhTM between 2011 and 2013 and 6 cases were excluded by central assessment. Sixty-one cases were included in the control arm. Serum samples on day 1 were obtained from 28 AE-IIPs patients out of 39 AE-IIPs patients in rhTM arm. Twenty-six AE-IIPs patients survived for more than 7 days and serum samples on day 8 were obtained from 22 patients out of the 26 patients.

Abbreviations: MPA, microscopic polyangiitis; AE, acute exacerbation; rhTM, recombinant human soluble thrombomodulin; IIPs, idiopathic interstitial pneumonias; FDP, fibrin degradation product.